Hospital:





First Name	Surname		Patient's postcode	
			-	
	_			
Date of Birth (dd/mm/yyyy)	Sex		Contact telephone number	
//	☐ Male ☐ Female ☐ Other			
Hospital MRN	Patient type		Indigenous Status	
Medicare number	□ Public □ Private □ Overseas □ Not known		 □ Aboriginal □ Torres Strait Islander □ Both Aboriginal and Torres Strait Islander □ Neither Aboriginal nor Torres Strait Islander □ Not known 	
Admission via ED of operating hospital		If transferred from another hospital		
 ☐ Yes ☐ No, transferred from another hospital ☐ No, in-patient fall ☐ Other/not known 		Name of transferring hospital: ED/Hospital arrival date/time//hrs (transferring hospital)		
ED/Hospital Admission (operating h	osnital)	Record time using 24hr clock If an in-patient fracture (time using 24hr clock)		
EDMOSPILAL Admission (operating in	oopitaly	in an in-patient fracture (time using 2-in clock)		
Admission// hrs		Date / time of diagnosis / / :hrs		
Departure//(from ED)	:hrs Record time using 24hr clock	Record time using 24hr clock		
Usual Place of Residence		Type of ward admitted to		
 □ Private residence including retirement village □ Residential care facility □ Other □ Not known 		☐ Hip fracture unit /Orthopaedic ward / preferred ward ☐ Outlying ward ☐ HDU / CCU / ICU ☐ Other / not known		
Note: If holiday residence/respite care, document usual place of residence		Preadmission cognitive		
Walking ability pre-admission		status	Preoperative cogni	itive assessment
□ Usually walks without walking aids □ Usually walks with a stick or crutch □ Usually walks with two aids or frame □ Usually uses a wheel chair/ bed bound □ Not known Note: if a person has different levels of mobility on different surfaces then		□ Normal cognition □ Impaired cognition or kn dementia □ Not known	□ Not assessed □ Assessed and no □ Assessed and ab □ Not known Note: cognitive assessi	normal or impaired
record the level of most assistance	•	Data Managara	validated tool	·
Pain Assessment	hin 30 minutes of ED	Pain Management		
□ Documented assessment of pain within 30 minutes of ED presentation □ Documented assessment of pain greater than 30 minutes of ED presentation □ Pain assessment not documented or not done		 □ Analgesia given within 30 minutes of ED presentation □ Analgesia given more than 30 minutes after ED presentation □ Analgesia provided by paramedics □ Analgesia not required □ Not known 		
Bone protection medication at admission		Clinical Frailty Preinjury Status		
□ No bone protection medication □ Yes, calcium and/or vitamin D only □ Yes, bisphosphonate (oral or IV) denosumab or teriparatide (with or without calcium and/or vitamin D) □ Not known		□ 1 Very fit □ 6 Moderately frail □ 2 Well □ 7 Severely frail □ 3 Well with treated comorbid conditions □ 8 Very severely frail □ 4 Vulnerable □ 9 Terminally ill □ 5 Mildly frail □ Not known		
Pre-operative medical assessment		Side of fracture		
□ No assessment conducted □ Geriatrician / geriatric team □ Physician / physician team □ GP □ Specialist nurse □ Not known		☐ Left ☐ Right If bilateral – complete a separate record for each fracture		
This is in addition to preoperative anaesthetic and orthopaedic review				
Atypical fracture Not a pathological or atypical fracture Pathological fracture Atypical fracture See data dictionary if uncertain of definitions		Type of fracture □ Intracapsular – undisplaced / impacted □ Intracapsular - displaced □ Per / intertrochanteric □ Subtrochanteric Note: Basal/basicervical #s are to be classed as per/intertrochanteric		

Did the patient undergo surgery	Date & time of primary surgery		
☐ Yes			
☐ No - surgical fixation not clinically indicated	/ / hrs		
☐ No - patient for palliation			
☐ No - other reason	Record time using 24hr clock		
Reason if delay > 48 hours	Anaesthesia		
☐ No delay - surgery < 48 hrs			
☐ Delayed due to patient deemed medically unfit	☐ General anaesthetic		
☐ Delayed due to issues with anticoagulation	☐ Spinal / regional anaesthesia		
☐ Delayed due to theatre availability	☐ General and spinal/regional anaesthesia		
☐ Delayed due to surgeon availability	□ Other – state		
☐ Delayed due to delayed diagnosis of hip fracture	□ Not known		
☐ Other type of delay (state reason)			
□ Not known			
Note: Delay is calculated from time of presentation to ED of the first			
hospital or diagnosis of hip fracture for those with a fracture from an in-			
patient fall			
Analgesia (nerve block)	Consultant present during surgery		
☐ Nerve block administered preoperative (before arriving in OT)	□ No		
☐ Nerve block administered in OT	□ Yes		
□ Both	☐ Not known		
☐ Neither			
☐ Not known	Note: To record yes, consultant must be scrubbed and operating		
Operation Performed	ASA Grade		
☐ Cannulated screws (e.g. multiple screws)			
☐ Sliding hip screw	□ 1 □ 2 □ 3 □ 4 □ 5 □ unknown		
☐ Intramedullary nail – short			
☐ Intramedullary nail – long			
☐ Hemiarthroplasty – stem cemented			
☐ Hemiarthroplasty – stem uncemented			
☐ Total hip replacement – stem cemented			
☐ Total hip replacement – stem uncemented			
Other			
□ Not known			
Postoporativo weight hearing status	First day mobilisation		
Postoperative weight bearing status Unrestricted weight bearing	☐ Patient out of bed and given opportunity to start mobilising day 1 post		
□ Restricted / non weight bearing	surgery		
□ Not known	☐ Patient not given opportunity to start mobilising day 1 post surgery		
1 NOT KHOWIT	□ Not known		
New Pressure Injury of the Skin	Delirium assessment		
New Freedom mjury of the oran			
□ No □ Yes □ Not known	□ Not assessed		
□ No □ Yes □ Not known			
☐ No ☐ Yes ☐ Not known Note: Grade 2 + above during acute admission	□ Not assessed □ Assessed and not identified		
	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known		
Note: Grade 2 + above during acute admission	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician □// Bone protection medication at discharge from operating hospital		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment No	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician □ / / Bone protection medication at discharge from operating hospital □ No bone protection medication		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment No Performed during admission	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician ■ / / Bone protection medication at discharge from operating hospital □ No bone protection medication □ Yes, calcium and/or vitamin D only		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment No Performed during admission Awaits falls clinic assessment	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician □ / / Bone protection medication at discharge from operating hospital □ No bone protection medication □ Yes, calcium and/or vitamin D only □ Yes, bisphosphonate (oral or IV) denosumab or teriparatide		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment No Performed during admission Awaits falls clinic assessment Further intervention not appropriate	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician □ — / — — / — — — Bone protection medication at discharge from operating hospital □ No bone protection medication □ Yes, calcium and/or vitamin D only □ Yes, bisphosphonate (oral or IV) denosumab or teriparatide (with or without calcium and/or vitamin D)		
Note: Grade 2 + above during acute admission Clinical malnutrition assessment Not done Malnourished Not malnourished Not known Assessed by Geriatrician in acute phase of care No Yes No geriatric medicine service available Not known Specialist falls assessment No Performed during admission Awaits falls clinic assessment	□ Not assessed □ Assessed and not identified □ Assessed and identified □ Not known Note: assessment of delirium requires use of a validated tool First day walking □ No □ Yes □ Not known Date initially assessed by Geriatrician □ / / Bone protection medication at discharge from operating hospital □ No bone protection medication □ Yes, calcium and/or vitamin D only □ Yes, bisphosphonate (oral or IV) denosumab or teriparatide		

Discharge

Date of discharge from acute / orthopaedic ward	Discharge destination from acute / orthopaedic ward	
/	□ Private residence (including retirement village) □ Residential care facility □ Rehabilitation unit - public □ Rehabilitation unit - private □ Other hospital / ward / speciality department □ Deceased □ Other □ Not known	
Date of final discharge from hospital if known	Discharge destination from hospital if known	
/	 □ Private residence (including retirement village) □ Residential aged care facility □ Deceased □ Other □ Not known 	

Follow Up 120 days

	120 days		
Follow up date	Note: record date that follow up was completed		
Alive at 120 days	☐ Yes Confirm date of final discharge from hospital system//		
Residential status	□ Private residence (including unit in retirement village) □ Residential aged care facility □ Rehabilitation unit - public □ Rehabilitation unit - private □ Other hospital / ward / speciality department □ Deceased □ Other □ Not known		
Walking ability	□ Usually walks without walking aids □ Usually walks with a stick or crutch □ Usually walks with two aids or frame □ Usually uses a wheel chair/ bed bound □ Not known		
Bone protection	□ No bone protection medication □ Yes - Calcium and/or vitamin D only □ Yes - Bisphosphonate (oral or IV) denosumab or teriparatide (with or without calcium and/or vitamin D) □ Not known		
Re-operation within120 days	□ No reoperation □ Reduction of dislocated prosthesis □ Washout or debridement □ Implant removal □ Revision of internal fixation □ Conversion to Hemiarthroplasty □ Conversion to THR □ Excision arthroplasty □ Revision arthroplasty □ Not relevant □ Not known		



Health Questionnaire


English version for Australia

Under each heading, please tick the ONE box that best describes your health TODAY. **MOBILITY** I have no problems with walking around I have slight problems with walking around I have moderate problems with walking around I have severe problems with walking around I am unable to walk around PERSONAL CARE I have no problems with washing or dressing myself I have slight problems with washing or dressing myself I have moderate problems with washing or dressing myself I have severe problems with washing or dressing myself I am unable to wash or dress myself **USUAL ACTIVITIES** (e.g. work, study, housework, family or leisure activities) I have no problems doing my usual activities I have slight problems doing my usual activities I have moderate problems doing my usual activities I have severe problems doing my usual activities I am unable to do my usual activities **PAIN / DISCOMFORT** I have no pain or discomfort I have slight pain or discomfort I have moderate pain or discomfort I have severe pain or discomfort I have extreme pain or discomfort **ANXIETY / DEPRESSION** I am not anxious or depressed I am slightly anxious or depressed I am moderately anxious or depressed I am severely anxious or depressed

I am extremely anxious or depressed

The best health you can imagine

100



30

25

20

15

10

5

The worst health

you can imagine

- We would like to know how good or bad your health is TODAY.
- This scale is numbered from 0 to 100.
- 100 means the <u>best</u> health you can imagine.
 0 means the <u>worst</u> health you can imagine.
- Mark an X on the scale to indicate how your health is TODAY.
- Now, please write the number you marked on the scale in the box below.

YOUR HEALTH TODAY =