

Data Dictionary
Version 14

October 2021

Australian and New Zealand Hip Fracture Registry

Background: A minimum data set (MDS) was created for the Australian and New Zealand (ANZ) Hip Fracture Registry Steering Group. The purpose of the minimum dataset and registry is to support consistent, local collection of data across Australia and New Zealand to enable facilities to compare practice and outcomes for hip fracture care against national clinical guidelines and standards of care. The MDS has been developed to capture information relevant to ANZ Hip Fracture Guidelines and national Hip Fracture Care Clinical Care Standard and is comparable to the United Kingdom (UK) national hip fracture registry and other registries emerging across the world.

Purpose: The ultimate goal of the ANZ Hip Fracture Registry is to use data to improve performance and maximise outcomes for older people who fracture their hip by reducing mortality and morbidity, reducing rates of institutionalisation, maximising functional independence and preventing future fractures by monitoring secondary prevention interventions.

MDS development: The MDS has been reviewed by the ANZ Hip Fracture Registry Steering Group, which consists of representatives of key professional and consumer bodies from Australia and New Zealand: Australian and New Zealand Society for Geriatric Medicine (ANZSGM); Australian Orthopaedic Association (AOA); Australian and New Zealand College of Anaesthetists (ANZCA); Australasian College of Emergency Medicine (ACEM); New Zealand Orthopaedic Association (NZOA); Royal Australasian College of Surgeons (RACS); Royal Australasian College of Physicians (RACP); Australian and New Zealand Orthopaedic Nurses Association (ANZONA); Australasian Faculty of Rehabilitation Medicine (AFRM); Australian Physiotherapy Association (APA); Osteoporosis Australia (OA); and Osteoporosis New Zealand (ONZ). This version of the ANZHFR Data Dictionary includes data variables for both the Patient Level Audit (the Registry) and the Facility Level Audit (annual snapshot of hospital level processes and protocols).

The data variables collected in the MDS (Patient Level) are from six (6) key components of care and include: (1) Patient information; (2) Admission; (3) Assessment; (4) Treatment; (5) Discharge; and (7) 120 day follow-up. The data variables collected in the MDS (Facility Level) cover: (1) Hospital Information; (2) Model of Care; (3) Protocols and processes; (4) Beyond the acute hospital stay; (5) Other aspects of care.

Core and non-core data items

Core variables are those variables collected by all using the minimum dataset e.g. date and time of admission, or type of hip fracture, and will be uploaded to the ANZ Hip Fracture Registry (ANZHFR). A number of these items will be considered mandatory for the purposes of forming a meaningful registry. Non-core items are collected at a local level and are held either locally or on the central server, or are generated automatically at a central level using data uploaded.

Review: The MDS will be reviewed annually by the ANZHFR Steering Group. It is anticipated that any new item to be added must be presented with a clear case for the benefits of adding it. Equally the Committee will be charged with removing redundant items which are felt not to add value either at a facility or central level.

Patient Inclusion: A person aged 50 years and older, who has been admitted to a participating hospital with an acutely fractured hip from a minimal or low trauma injury, and who undergoes either surgical or non-surgical management of the hip fracture.

Version history:

Version	Description of Change	Author	Date Changed	Status
1.0	Draft	Rebecca Mitchell	July 2012	Rough draft
2.0	Draft	Rebecca Mitchell	6 Dec 2012	Draft
3.0	Draft	Rebecca Mitchell	8 Jan 2013	Draft
4.0	Draft	Rebecca Mitchell	3 May 2013	Draft
5.0	Draft	Rebecca Mitchell	31 May 2013	Draft
6.0	Penultimate	Rebecca Mitchell	21 June 2013	Draft
7.0	Final	Rebecca Mitchell	5 July 2013	Final
8.0	Revised to ensure data capture consistent with Guideline recommendations	Jacqui Close	4 Dec 2013	Final
8.1	Revised to ensure data capture consistent with Guideline recommendations and the requirement to capture identifying data for follow up and data linkage	Jacqui Close	11 Dec 2013	Final
9.0	Review by the Steering Group against the 2014 ANZ Guidelines for Hip Fracture Care and the 2016 ACSQHC Hip Fracture Care Clinical Care Standard and Indicators; incorporation of definitions for the Facility Level Audit variables	Elizabeth Armstrong	August 2016	Draft
9.1	Revision with Steering Group and Data Committee feedback	Elizabeth Armstrong	September 2016	Final Draft
10.0	Review by the Steering Group to incorporate feedback from participating sites and ensure data dictionary continues to be fit for purpose	Steering Group	August 2017	Draft
10.1	Revision with Data Committee feedback	Data Committee	October 2017	Draft
10.2	Revision with Data Committee feedback	Data Committee	October 2017	Final Draft
11	Annual Steering Group review to ensure data dictionary continues to be fit for purpose	Data Committee	October 2018	Final
12	Annual Steering Group review to ensure data dictionary continues to be fit for purpose	Data Committee	October 2019	Final Draft
13	Annual review of the dataset	Data Committee	October 2020	Final
14	Annual review of the dataset	Data Committee	October 2021	Draft

List of Data VariablesANZHFR Patient Level Audit	
Section 1 Patient information	7
Unique identifier	7
Australian and New Zealand jurisdiction	7
First name of patient	8
Surname of patient	8
Hospital MRN / URN / event number	8
Contact telephone number for patient	9
Date of birth	
Age derived	9
Sex of person	10
Australian Indigenous status	10
NZ ethnic status	11
Patient's postcode	122
Medicare number (Australia) / National Health Index (New Zealand)	122
Patient type	13
Usual place of residence	
Statistical linkage key 581	14
Section 2 Admission	
Establishment identifier of operating hospital	
Admission via ED of operating hospital	
Transfer hospital	
ED / hospital arrival date (transfer hospital)	
ED arrival time (transfer hospital)	
ED / other ward arrival date (operating hospital)	
ED / other ward arrival time (operating hospital)	
ED departure date (operating hospital)	
ED departure time (operating hospital)	
In-patient fracture date	
In-patient fracture time	
Pain assessment	20
Pain management	20
Ward type	21
Section 3 Assessment	
Pre-admission walking ability	
Pre-operative cognitive assessment	22
Pre-admission cognitive status	
Bone protection medication at admission	23
Pre-operative medical assessment	
Side of hip fracture	
Atypical fracture	25
Type of fracture	25
Surgical repair	26
ASA grade	26
Clinical Frailty Scale.	
Section 4 Treatment	
Date of surgery for hip fracture	28
Time of surgery for hip fracture	28
Surgery delay	
Surgery delay other text	
Type of anaesthesia	
Analgesia - nerve block	
Consultant surgeon present	
Type of operation performed	
Full weight bear	
First day mobilisation	

New pressure injuries of the skin	
•	
Assessed by geriatric medicine	
Geriatric medicine assessment date	
Specialist falls assessment	
Bone protection medication at discharge from acute hospital	
Delirium assessment	38
Clinical malnutrition assessment	39
First day walking	40
Section 5 Discharge	41
Discharge date from acute ward	41
Discharge destination from acute orthopaedic episode	41
Discharge from hospital date	
Length of stay (operating hospital)	42
Length of stay (health system)	43
Discharge place of residence	
Section 7 120 day follow-up*	45
120 day follow-up date	
Survival at 120 days post-surgery	45
Date health system discharge at 120 day follow-up	45
Place of residence at 120 day follow-up	46
Post-admission walking ability at 120 day follow-up	47
Bone protection medication at 120 day follow-up	48
Re-operation within 120 day follow-up	48
Preliminary date of death	49
Final date of death	49
Underlying cause of death	50
Other causes of death	50
ANZHFR Facility Level Audit	53
Section 8 Hospital information	53
Major trauma centre	53
Hip fractures	
Hip fractures Section 9 Model of care	53
·	53
Section 9 Model of care	53 54 54
Section 9 Model of care Orthogeriatric service	53 54 54
Section 9 Model of care Orthogeriatric service Model of care	53545555
Section 9 Model of care Orthogeriatric service Model of care Section 10 Protocols and processes	5354555556
Section 9 Model of care	
Section 9 Model of care Orthogeriatric service Model of care Section 10 Protocols and processes Imaging protocol Hip fracture pathway	
Section 9 Model of care Orthogeriatric service Model of care Section 10 Protocols and processes Imaging protocol Hip fracture pathway Venous thromboembolism protocol	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service Model of care Section 10 Protocols and processes Imaging protocol Hip fracture pathway Venous thromboembolism protocol Pain protocol Planned theatre list	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service	
Section 9 Model of care Orthogeriatric service Model of care Section 10 Protocols and processes Imaging protocol Hip fracture pathway Venous thromboembolism protocol Pain protocol Planned theatre list Anaesthesia Nerve block for pain pre-surgery Nerve block for pain post-surgery Therapy access Section 11 Beyond the Acute Hospital Stay Information on treatment and care Inpatient rehabilitation Home-based rehabilitation Injury prevention Falls clinic	
Section 9 Model of care Orthogeriatric service	53 54 54 55 56 56 56 57 57 58 58 59 60 61 61 61 62 63 63
Section 9 Model of care	53 54 54 55 56 56 56 57 57 58 58 60 61 61 61 62 63 63
Section 9 Model of care	
Section 9 Model of care	53 54 55 55 56 56 57 57 58 58 59 60 61 61 61 62 63 63 64 64

Service provision plan details	66
Service provision barriers	67
Service provision barrier details	
Hip fracture data collector	67
Hip fracture data entry	
Time taken data collection and entry	
Time taken 120-days	
Sprint audits	
Impact of COVID-19	

ANZHFR Patient Level Audit

Section 1 Patient information

Variable Number 1.01

Variable Unique identifier

Variable Name

Definition A consecutive number allocated to each record of a hip fracture

Justification To allow for the identification of records

Format 10 digit numeric

Status Non-core (created centrally)

Coding Source
Coding Frame

DD Comments This is the unique record number used to identify each record

Variable Number 1.02

Variable Australian and New Zealand jurisdiction

Variable Name Area

Definition The Australian or New Zealand jurisdiction of the hospital

Justification To enable the identification of hospitals in Australian and New Zealand jurisdictions

Format 2 digit numeric

Status Non-core (created centrally)

Coding Source Adapted from the National Health Data Dictionary, Version 15 (METeOR identifier

269941)

Coding Frame 1 New South Wales

2 Victoria3 Queensland4 South Australia5 Western Australia

6 Tasmania

7 Northern Territory

8 Australian Capital Territory

9 Other Territories (Cocos Keeling Islands, Christmas Island and Jervis Bay Territory)

10 New Zealand

DD Comments The order used here is the standard for the Australian Bureau of Statistics (ABS).

Variable First name of patient

Variable Name Name

Definition First name of the patient

Justification To allow for checking of duplicate entries for the one person and to contact the

patient for the 120 day follow-up

Format Character Status Core

Coding SourceTo allow for checking of duplicate entries for the one person as well as the ability to

follow up patient including future data linkage

Coding Frame Character string

DD CommentsThe format should be the same as that indicated by the person (for example written

on a form) or in the same format as that printed on an identification card, such as

Medicare card, to ensure consistent collection of name data

Variable Number 1.04

Variable Surname of patient

Variable Name Surname

Definition Surname of the patient

Justification To allow for checking of duplicate entries for the one person as well as the ability to

follow up patient including future data linkage

Format Character Status Core

Coding Source Coding Frame

DD CommentsThe format should be the same as that indicated by the person (for example written

on a form) or in the same format as that printed on an identification card, such as

Medicare card, to ensure consistent collection of name data

Variable Number 1.05

Variable Hospital MRN / URN / event number

Variable Name MRN

Definition Hospital Medical Record Number

Justification Unique person-identifier for each patient in each hospital and contributes to

collection of information on follow up e.g. re-operation

Format String XXXXXX[X(14)]

Status Core

Coding Source Coding Frame

DD Comments Key field: must be entered to create a patient record. Individual hospitals use their

own alphabetic, numeric, or alphanumeric coding systems. With the eventual move to E-Health in Australia, each patient will have a unique id nation-wide. Note: Western Australia uses URN. New Zealand to record the hospital event number.

Variable Contact telephone number for patient

Variable Name phone

DefinitionContact telephone number of the patient**Justification**To contact the patient for the 120 day follow-up

Format 10 digit numeric

Status Core

Coding Source Coding Frame

DD CommentsOnly record one telephone number. This should be the best land line telephone or

mobile phone number to contact the patient for the 120 day follow-up. Record the prefix plus telephone number without punctuation, for example, 08 8226 6000 or

0417 123456.

Variable Number 1.07

Variable Date of birth

Variable Name DOB

Definition Date of birth of the patient

Justification Basic demographic details. Required for probabilistic data linkage

Format 8 digit, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 287007)

Coding Frame DD/MM/YYYY

DD Comments Key field Australia: must be entered to create a patient record. Only include people

who are 50 years and older at the time of their hip fracture admission. Date not

known is recorded as: 01011900.

Variable Number 1.08

Variable Age derived

Variable Name Age

Definition Age of the patient in (completed) years at admission

Justification Basic demographic details

Format 3 digit, N[NN]

Status Non-core (created centrally)

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 303794)

Coding Frame 999 Unknown/Not stated

DD Comments If age (or date of birth) is unknown or not stated, and cannot be estimated, use

Code 999. Age to be calculated automatically from Date of Birth and ED/hospital arrival date (operating hospital) or ED/hospital arrival date (transfer hospital) for

patients transferred to an operating hospital

Variable Sex of person

Variable Name Sex

Definition Sex of the patient

Justification Basic demographic details

Format 1 digit numeric

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame 1 Male

2 Female

3 Intersex or indeterminate

9 Not stated / inadequately described

DD Comments Key field: must be entered to create a patient record.

Variable Number 1.10

Variable Australian Indigenous status

Variable Name Indig

Definition Was the patient of Aboriginal or Torres Strait Islander origin?

Justification Basic demographic details

Format 1 digit numeric, N

Status Core

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 291036)

Coding Frame 1 Aboriginal but not Torres Strait Islander origin

2 Torres Strait Islander but not Aboriginal origin3 Both Aboriginal and Torres Strait Islander origin4 Neither Aboriginal or Torres Strait Islander origin

9 Not stated / inadequately described

DD Comments An Aboriginal or Torres Strait Islander is a person of Aboriginal or Torres Strait

Islander descent who identifies as an Aboriginal or Torres Strait Islander and is accepted as such by the community in which he or she lives. Collected Australia only

Variable NZ ethnic status

Variable Name ethnic

Definition Which ethnic group or groups does the patient belong to?

Justification Basic demographic details

Format 1 digit numeric

Status Core

Coding Source Ministry of Health. 2017. HISO 10001:2017 Ethnicity Data Protocols. Wellington:

Ministry of Health

Coding Frame 10 European

11 New Zealand European

12 Other European

21 Māori

30 Pacific peoples not further defined

31 Samoan

32 Cook Islands Māori

33 Tongan34 Niuean35 Tokelauan36 Fijian

37 Other Pacific Peoples 40 Asian not further defined

41 Southeast Asian

42 Chinese 43 Indian 44 Other Asian 51 Middle Eastern 52 Latin American 53 African

61 Other Ethnicity94 Don't Know95 Refused to answer97 Response unidentifiable

99 Not stated

DD Comments Patients should be asked to self-identify their ethnicity by asking them 'Which

ethnic group or groups do you belong to?' For many patients it will not be possible to ask them this during their hospital admission. Therefore, the ethnicity that is recorded in the NZ hospital system should be used. The accuracy of ethnic group(s) can then be clarified at the 120 day follow up phone call. The collector must not limit the number of ethnicities given. Decisions around reporting of ethnic groups will be made in consultation with NZOA Nga Rata Koiwi representative on the NZIMC (New Zealand Implementation Committee). Collected New Zealand only.

Variable Patient's postcode

Variable Name Apcode

Definition What was the postcode of the suburb of the usual residence of the patient?

JustificationBasic demographic detailsFormat4 digit numeric, {NNNN}

Status Core

Coding Source Australia Post or New Zealand Post websites (<u>www.auspost.com.au or</u>

www.nzpost.co.nz) provide up-to-date postcodes and localities

Coding Frame 1000 No fixed abode

9998 Overseas

9999 Postcode not known

DD Comments Use a valid Australian or New Zealand postcode

Variable Number 1.13

Variable Medicare number (Australia) / National Health Index (New Zealand)

Variable Name Medicare

Definition Patient's Medicare number

JustificationTo allow for checking of duplicate entries for the one person and for multiple

admissions

Format Characters, N(11)

Status Core

Coding Source Coding Frame

DD Comments Enter the full Medicare number for an individual (i.e. family number

plus person individual reference number).

Key field New Zealand: must be entered to create a patient record. New Zealand will provide the National Health Index (NHI) which is a unique number assigned to every person who uses health and disability services in New Zealand. New Zealand will

use this variable as the main mechanism to identify each patient.

Variable Patient type

Variable Name ptype

Definition Payment status

JustificationTo identify the source of revenue received by a health industry relevant organisation

Format 3 digit numeric

Status Core

Coding Source Adapted from the National Health Data Dictionary, Version 15

Coding Frame 1 Public

2 Private3 Overseas9 Not known

DD Comments For New Zealand all surgery for hip fractures takes place in the public sector. There

will be the occasional patient from overseas and this should be noted accordingly.

In Australia, private sector patients include those with treatment funded by: private health insurance, workers' compensation insurance, motor vehicle third party insurance, other compensation (e.g. Public liability, common law, medical negligence), private households (i.e. self-funded and out-of-pocket expenditure), non-profit institutions serving households, corporations (other than health

insurance), other private sector revenue

In Australia, public sector patients include those with treatment funded by: Medicare, Pharmaceutical Benefits Scheme and Repatriation Pharmaceutical Benefits Scheme, Department of Veterans' Affairs, National Health and Medical Research Council, Australian Health Care Agreements, other Special Purpose payments, Other Australian Government Departments, State/Territory non-health

departments, or other public sector revenue

Variable Usual place of residence

Variable Name uresidence

Definition What is the usual place of residence of the patient?

Justification Type of accommodation before and after admission are collected to compare where

the patient has come from (what was their usual accommodation) and where they are going to (what will become their usual accommodation). This is an indicator of

patient outcome.

Format 1 digit numeric

Status Core

Coding Source Adapted from the Australasian Rehabilitation Outcomes Centre Inpatient Dataset,

Version 3.0; NSW SNAP Data Collection, Version 4.0

Coding Frame 1 Private residence (including unit in retirement village)

2 Residential aged care facility

3 Other 4 Not known

DD Comments Record the patient's usual accommodation type at admission.

Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services

in Australia and private hospitals or rest homes in New Zealand.

If the patient lives with a relative or in a community group home or boarding house

code 'private residence'.

If the patient was admitted from respite care, record their usual place of residence

when not in respite care.

Variable Number 1.16

Variable Statistical linkage key 581

Variable Name slk581

Definition A specific code (key) that can be used to bring together two or more records

belonging to the same individual. It is represented by a code consisting of characters

from the person's surname, first name, date of birth and gender.

Justification Brings together data from different sources to enable greater understanding of the

utilisation of health care and/or services. Clinical quality registries should have the capacity to enhance their value through the use of linkage to other datasets (Australian Commission on Safety and Quality in Health Care Framework for

Australian Clinical Quality Registries 2014)

Format 14 Characters XXXXXDDMMYYYYN

Status Core (created centrally)

Coding Source National Health Data Dictionary, Version 16 (METeOR identifier 349895)

Coding Frame

DD Comments It is represented by a code consisting of the second, third and fifth characters of a

person's family name, the second and third letters of the person's given name, the day, month and year when the person was born and the sex of the person, concatenated in that order. In Australia, the linkage key is designed to make it possible to count number of clients and services they received, without counting the same client more than once. It can also be used for linking to other related data collections. This may be useful for New Zealand, although the NHI is usually the best and only identifier used for data matching in New Zealand.

Section 2 Admission

Variable Number 2.01

Variable Establishment identifier of operating hospital

Variable Name Ahoscode

Definition Name of the operating hospital where the patient received surgery for the hip

fracture

Justification To allow for the identification of the establishment for benchmarking and

comparison purposes

Format Character Status Core

Coding Source Coding Frame

DD CommentsNote: For data analysis each hospital will have to be given a unique number

Variable Number 2.02

Variable Admission via ED of operating hospital

Variable Name EDadmit

Definition Did the patient present directly to the ED of the operating hospital?

Justification Ability to monitor the time spent in ED.

Format 1 digit Status Core

Coding Source

Coding Frame 1 Yes

2 No - transferred from another hospital

3 No - in-patient fall 9 Other / not known

DD Comments If the patient was admitted via the ED of the operating hospital, information on the

date and time that the patient arrived and left the ED of the operating hospital will

be recorded.

Variable **Transfer hospital**

Variable Name Athoscode

Definition Name of the hospital where the patient first presented and was diagnosed with a

hip fracture

Justification To allow for the identification of the establishment for benchmarking and

comparison purposes

Format Character Status Core

Coding Source

Coding Frame Not transferred

If transferred enter hospital name of first transfer hospital

DD Comments If the patient has not been transferred, this field will be hidden and will not be

available.. Note: For data analysis, each hospital will be given a unique number.

If patient is not transferred, data variables 2.04 and 2.05 regarding transfer

date/time will be hidden and will not be available.

Variable Number 2.04

Variable ED / hospital arrival date (transfer hospital)

Variable Name tarrdate

Definition Date on which the patient presented to the transferring hospital with a hip fracture **Justification**

To enable the identification of the date of arrival in transferring hospital. Will allow

for quantification of true time to surgery and overall LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments If the patient is transferred several times, this should be the hospital where the

patient first presented with the hip fracture.

If the presenting hospital has no ED or the patient wasn't admitted through ED, state the date presenting to the transferring hospital with a hip fracture. If the hip fracture occurred as an in-patient, record the date the fracture was

diagnosed.

Note: 00000000 indicates that the patient did not present through the ED and 9999998 indicates that patient was not transferred (i.e. not relevant) and 01011900 indicates that the date was not known. To be used in the calculation of

time to surgery and total LOS in the health system for the care episode.

Variable ED arrival time (transfer hospital)

Variable Name tarrtime

Definition Time at which the patient arrived in the ED of the transferring hospital

Justification To enable the identification of the time of arrival in the ED

Format 4 digit Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame hh:mm

DD Comments Time is recorded using the 24 hour clock.

If the patient is transferred several times, this should be the hospital where the

patient first presented with a hip fracture.

If the presenting hospital has no ED or the patient wasn't admitted through ED, state the time presenting to the transferring hospital with a hip fracture.

If the hip fracture occurred as an in-patient, record the time the fracture was

diagnosed.

Note: 0000 indicates that the patient did not present through the ED, 9998 indicates that patient was not transferred (i.e. not relevant), and 9999 indicates that time was not known. To be used in the calculation of total LOS in the health system for the

care episode.

Variable Number 2.06

Variable ED / other ward arrival date (operating hospital)

Variable Name arrdate

DefinitionDate on which the patient arrived in the ED / other ward of the operating hospital **Justification**To enable calculation of age at presentation, time spent in ED, time to surgery and

LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments If the patient was not admitted through the ED but was transferred from another

hospital and admitted directly to a ward of the operating hospital, state the date

admitted to the ward of the operating hospital. $% \label{eq:controlled}$

If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded. The Australian National Emergency Access Target (NEAT) aims that by 2015, 90% of patients will leave the ED within 4 hours — either by discharge, being admitted to

hospital or through transfer to another hospital for treatment

(http://www.ecinsw.com.au/node/128). For New Zealand patients are expected to

be discharged or admitted to hospital within 6 hours.

Variable ED / other ward arrival time (operating hospital)

Variable Name arrtime

Definition Time at which the patient arrived at the ED / other ward of the operating hospital

Justification To enable calculation of time spent in ED, time to surgery and LOS

Format 5 digit Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame hh:mm

DD Comments Time is recorded using the 24 hour clock.

If the patient was not admitted through the ED but was transferred from another hospital and admitted directly to a ward of the operating hospital, state the time

admitted to the ward of the operating hospital.

Note: 9999= time not known. If the patient was admitted via the ED of the operating

hospital, information on the date and time that the patient left the ED of the

operating hospital will be recorded

Variable Number 2.08

Variable ED departure date (operating hospital)

Variable Name depdate

Definition Date on which the patient departed from the ED of the operating hospital

Justification To enable calculation of time spent in ED, time to surgery and LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments Note: 01011900 indicates that the patient did not present through the ED. If the

patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded

Variable Number 2.09

Variable ED departure time (operating hospital)

Variable Name deptime

Definition Time at which the patient departed from the ED of the operating hospital

Justification To enable calculation of time spent in ED, time to surgery and LOS

Format 4 digit Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame hh:mm

DD Comments Time is recorded using the 24 hour clock.

Note: 9999= time not known. If the patient was admitted via the ED of the operating

hospital, information on the date and time that the patient left the ED of the

operating hospital will be recorded.

Variable In-patient fracture date

Variable Name admdateop

Definition Date on which the admitted patient commences the episode of care at the

operating hospital with radiological-confirmed diagnosis of hip fracture

Justification To enable the identification of the date of hip fracture occurring as an in-patient and

calculation of time to surgery and LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments Note: 01011900 = date not known

Fractures sustained while on leave from an existing hospital admission are not classified as inpatient fractures. They are recorded as a new event and date and

time of presentation are recorded at 2.06 and 2.07.

Variable Number 2.11

Variable In-patient fracture time

Variable Name admtimeop

Definition 24-hour time at which the admitted patient commences the episode of care at the

operating hospital with radiological-confirmed diagnosis of hip fracture

Justification To enable the identification of the time of hip fracture occurring as an in-patient

and calculation of time to surgery and LOS

Format 4 digit Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame hh:mm

DD Comments Time is recorded using the 24 hour clock. Note: 9999 = time not known

Variable Pain assessment

Variable Name painassess

Definition Did the patient have a documented assessment of pain within 30 minutes of

presentation to the emergency department

Justification Acute pain associated with the hip fracture can have adverse effects on outcome.

Hip Fracture Care Clinical Care Standard Indicator 2b.

Format 1 digit Status Core

Coding Source

Coding Frame 1 Documented assessment of pain within 30 minutes of ED presentation

2 Documented assessment of pain greater than 30 minutes of ED presentation

3 Pain assessment not documented or not done

9 Not known

DD Comments A pain assessment is any qualitative or quantitative assessment of pain recorded in

the notes.

A standardised pain assessment system should be used that specifically addresses the needs of patients with cognitive impairment and those unable to communicate pain. Time to pain assessment in the ED to be identified from clinical notes. Time is calculated from date and time of presentation to the emergency department of the

first hospital.

Variable Number 2.13

Variable Pain management

Variable Name painmanage

Definition Did the patient receive analgesia within 30 minutes of presentation to the

emergency department?

Justification Acute pain associated with the hip fracture can have adverse effects on outcome.

Hip Fracture Care Clinical Care Standard Indicator 2b.

Format 1 digit Status Core

Coding Source

Coding Frame 1 Analgesia given within 30 minutes of ED presentation

2 Analgesia given more than 30 minutes after ED presentation
3 Analgesia not required – already provided by paramedics
4 Analgesia not required – no pain documented on assessment

9 Not known

DD CommentsTime to analgesia in the ED to be identified from clinical notes. Time is calculated

from date and time of presentation to the emergency department of the first

hospital.

Variable Ward type

Variable Name ward

Definition What type of ward was the patient admitted to from ED?

Justification To enable the identification of the ward where the patient commenced their

episode of care

Format 1 digit Status Core

Coding Source

Coding Frame 1 Hip fracture unit/Orthopaedic ward/ preferred ward

2 Outlying ward 3 HDU / ICU / CCU 9 Other/ not known

DD Comments HDU refers to High Dependency Unit. ICU refers to Intensive Care Unit. CCU refers

to Coronary Care Unit.

An outlying ward refers to a ward not clinically appropriate to meet the patient's

current needs.

Section 3 Assessment

Variable Number 3.01

Variable Pre-admission walking ability

Variable Name walk

Definition What was the patient's walking ability pre-admission?

Justification To enable the identification of the mobility status pre-admission

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 Usually walks without walking aids

2 Usually walks with either a stick or crutch

3 Usually walks with two aids or frame (with or without assistance of a person)

4 Usually uses a wheelchair / bed bound

9 Not known

DD Comments If a person has different levels of mobility on different surfaces then record the level

of most assistance. For example, inside their residence a person usually walks without a walking aid but when outside the residence the person usually walks with

a frame, then the level of mobility recorded is option 3.

Variable Number 3.02

Variable Pre-operative cognitive assessment

Variable Name cogassess

Definition Following admission to hospital, cognitive status is assessed prior to surgery using a

validated tool and recorded in the medical record

Justification Hip fracture patients are at high risk of having an existing cognitive impairment or

developing delirium. Cognitive impairment and delirium in these patients is associated with increased morbidity and mortality, and a decrease in rehabilitation

potential and return to pre-fracture functioning.

Care at Presentation Hip Fracture Care Clinical Care Standard Indicator 1b.

Format 1 digit Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 Not assessed

2 Assessed and normal

3 Assessed and abnormal or impaired

9 Not known

DD Comments Cognitive assessment requires the use of a validated tool. Some validated tools for

assessing cognitive function include:

Abbreviated Mental Test Score (AMTS) (Hodkinson 1972)

Standardised Mini-Mental State Examination (SMMSE) (Molloy & Standish
 Standardised Mini-Mental State Examination (SMMSE) (Molloy & Standish

1997)

Modified Mini Mental State Exam (3MS) (Teng & Chui 1987)

 General Practitioner's Assessment of Cognition (GPCOG) (Brodaty et al. 2002)

The 4AT (Bellelli et al. 2014)

 Other tools, such as the Rowland Universal Dementia Assessment Scale (RUDAS) (Storey et al. 2004) and the Kimberly Indigenous Cognitive Assessment (KICA) (LoGiudice et al. 2006), may be more appropriate for

some people from culturally and linguistically diverse groups

Variable Pre-admission cognitive status

Variable Name cogstat

Definition What was the cognitive status of the patient prior to admission?

Justification To enable the identification of the cognitive status of the patient prior to admission.

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 1 Normal cognition

2 Impaired cognition or known dementia

9 Not known

DD Comments Normal cognition refers to 'no history of cognitive impairment or dementia'. Impaired

cognition or known dementia refers to a 'loss of cognitive ability and/or a decline in memory or other thinking skills severe enough to reduce a person's ability to perform

everyday activities' (Alzheimer's Association).

Variable Number 3.06

Variable Bone protection medication at admission

Variable Name bonemed

Definition Was the patient taking bone protection medication prior to sustaining the hip

fracture?

Justification Ability to monitor use of bone protection medication prior to hip fracture

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No bone protection medication

1 Yes - calcium and/or vitamin D only

2 Yes - bisphosphonates, denosumab, romosozumab or teriparitide (with or without

calcium and/or vitamin D)

9 Not known

DD Comments Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or

one alpha).

Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate,

Zoledronate, Pamidronate.

Variable Pre-operative medical assessment

Variable Name passess

Definition Who conducted the pre-operative medical assessment apart from anaesthetic and

orthopaedic review?

Justification To determine level of pre-operative medical assessment. Hip Fracture Care Clinical

Care Standard Indicator 3a.

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No assessment conducted

1 Geriatrician / Geriatric Team 2 Physician / Physician Team

3 GP

4 Specialist nurse 9 Not known

DD Comments The pre-operative assessment is conducted in addition to an anaesthetic review and

orthopaedic assessment. If the pre-operative assessment is conducted by a number of assessment team members, select the highest numerical option in the coding frame drop down list eg. the highest numerical option to select is '1' geriatrician.

Variable Number 3.08

Variable Side of hip fracture

Variable Name side

Definition What was the side of the patient's hip fracture?

Justification To enable the identification of the side of the hip fracture

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 1 Left

2 Right

DD Comments Key field: must be entered to create a patient record.

If the patient has bilateral hip fractures, a separate record should be created for

each fracture.

Variable Atypical fracture

Variable Name afracture

Definition Was the type of the patient's hip fracture either pathological or atypical?

Justification To enable the identification of fractures which are not consistent with the nature of

the injury

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 Not a pathological or atypical fracture

1 Pathological fracture 2 Atypical fracture

DD Comments A pathological fracture is considered to be a fracture that has occurred when a bone

breaks in an area that has been weakened by another disease process (except osteoporosis), such as a tumour, infection or an inherited bone disorder.

An atypical fracture is one where the radiologically observed fracture pattern is not

consistent with the mechanism of injury described and is not thought to be

attributable to a discrete underlying disease process

Variable Number 3.10

Variable Type of fracture

Variable Name ftype

Definition What was the type of the patient's hip fracture? **Justification** To enable the identification of the type of hip fracture

Format 1 digit numeric

Status Core

Coding SourceAdapted from the UK National Hip Fracture DatabaseCoding Frame1 Intracapsular undisplaced/impacted displaced

2 Intracapsular displaced 3 Per/intertrochanteric 4 Subtrochanteric

DD CommentsBasal/basicervical fractures are to the classified as per/intertrochanteric

Variable Surgical repair

Variable Name surg

Definition Did the patient undergo surgical repair of the hip fracture?

Justification To enable quantification of percentage patients undergoing surgery

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame

2 Yes

3 No – surgical fixation not clinically indicated

4 No – patient for palliation

5 No – other reason

DD Comments

Variable Number 3.12

Variable ASA grade

Variable Name asa

Definition What is the ASA grade for the patient?

Justification A marker of disease severity and operative risk and used for case-mix adjustment

Format 1 digit numeric

Status Core

Coding Source American Society of Anaesthesiologists

1 Healthy individual with no systemic disc

Coding Frame 1 Healthy individual with no systemic disease

2 Mild systemic disease not limiting activity

3 Severe systemic disease that limits activity but is not incapacitating 4 Incapacitating systemic disease which is constantly life threatening 5 Moribund not expected to survive 24 hours with or without surgery

9 Not known

DD Comments ASA grade is used in case-mix adjustment for outcome at 30 and 120 days post-

surgery

Variable Clinical Frailty Scale

Variable Name frailty

Definition What was the patient's pre-injury frailty status?

Justification To enable the identification of the patient's frailty status prior to their hip fracture as a

person's level of frailty impacts outcomes.

Format 2 digit numeric

Status Core

Coding Source Rockwood Clinical Frailty Scale

Coding Frame 1 Very Fit

2 Well

3 Well, with treated comorbid disease

4 Vulnerable 5 Mildly frail 6 Moderately frail 7 Severely frail 8 Very severely frail 9 Terminally ill 99 Not known

DD Comments

NOTE: the Clinical Frailty Scale applies to the person's usual status prior to the hip fracture. Where the person has dementia or delirium the information will need to be provided by an informant who knows the person well.

Coding Frame Definitions

1 **Very fit** - robust, active, energetic and well-motivated. Exercise regularly and are among the fittest for their age.

2 **Well** - without active disease symptoms but are less fit than category 1. Exercise occasionally.

3 **Well with treated comorbid disease** - disease symptoms are well controlled compared to category four. Not regularly active beyond routine walking.

4 **Vulnerable** - not dependent on others for daily help, but symptoms limit activities. Common complaint is being 'slowed up' or being tired during the day.

5 **Mildly frail** - more evident slowing, and need help in instrumental activities of daily living (e.g. heavy housework, medications, transportation, shopping, using the phone, managing finances, meal preparation).

6 **Moderately frail** - need help with both instrumental and non-instrumental activities of daily living. Includes mobility in bed, transferring on/off chairs, toilets and into/out of bed, walking, dressing, eating, toilet use, personal hygiene, bathing.

7 **Severely frail** - completely dependent on others for all activities of daily living for whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~ 6 months).

8 **Very severely frail** - completely dependent on others for all activities of daily living, approaching the end of life. Typically, they could not recover even from a minor illness. 9 **Terminally ill** - approaching the end of life. Applies to people with a life expectancy <6 months who are not otherwise evidently frail.

Section 4 Treatment

Variable Number 4.01

Variable Date of surgery for hip fracture

Variable Name sdate

Definition Date on which the surgery for the hip fracture takes place

Justification To enable the identification of the date of primary surgery. Hip Fracture Care Clinical

Care Standard Indicator 4a.

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments If there was no surgery, enter 00000000. Date not known is classified as: 01011900

Variable Number 4.02

Variable Time of surgery for hip fracture

Variable Name stime

Definition 24-hour time at which the surgery for the hip fracture commences. This time is

taken from the start of the anaesthetic process.

Justification To enable the identification of the start time of the primary surgery. Hip Fracture

Care Clinical Care Standard Indicator 4a.

Format 4 digit Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame hh:mm

DD Comments Time is recorded using the 24 hour clock

The time of surgery for the hip fracture is taken from the start of the anaesthetic

process. Unknown time is classified as: 9999.

Variable Surgery delay

Variable Name delay

Definition What was the primary reason for the delay if the delay was greater than 48 hours

from the time of arrival in the emergency department of the first hospital, or

diagnosis of a fracture if the fracture occurred as an in-patient?

Justification Ability to monitor time to surgery as a standard of care

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 No delay, surgery completed <48 hours

2 Delay due to patient deemed medically unfit3 Delay due to issues with anticoagulation

4 Delay due to theatre availability 5 Delay due to surgeon availability

6 Delay due to delayed diagnosis of hip fracture

7 Other type of delay (state reason)

9 Not known

DD Comments Delay is calculated from the time of presentation in the emergency department of

the first hospital.

A person is considered medically unfit if he/she have acute health-related issues which need to be stabilised/optimised or reversed prior to proceeding with

anaesthesia and a surgical procedure.

If there is more than one delay to surgery, choose the reason for the first delay.

Variable Number 4.04

Variable Surgery delay other text

Variable Name delay_txt

Definition What was the reason for the other delay, if the delay was greater than 48 hours

from the time of arrival in the emergency department?

Justification Ability to monitor time to surgery as a standard of care

Format Character Status Core

Coding Source Coding Frame DD Comments

Variable Type of anaesthesia

Variable Name anaesth

Definition What type of anaesthesia for the hip fracture surgery?

Justification Ability to monitor variation, post-operative complications and patient choice

Format 2 digit numeric

Status Core

Coding Source

Coding Frame 1 General anaesthesia

5 Spinal / regional anaesthesia

6 General and spinal/regional anaesthesia

97 Other 99 Not known

DD Comments CSE=Combined Spinal/Epidural. Recorded in anaesthetic chart

Variable Number 4.06

Variable Analgesia - nerve block

Variable Name analges

Definition Did the patient have a nerve block?

Justification Monitoring against Guideline recommendation

Format 2 digit numeric

Status Core

Coding Source

Coding Frame 1 Nerve block administered before arriving in OT

2 Nerve block administered in OT

3 Both 4 Neither

99 Not known

DD Comments

Variable Consultant surgeon present

Variable Name consult

Definition Was the consultant surgeon operating or assisting with the operation?

Justification Ability to monitor the impact of consultant surgeon presence on the quality and

safety of patient outcome

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

DD CommentsTo record yes, consultant must be scrubbed and operating. This variable can be

found by checking if the consultant surgeon is recorded on the operation sheet

Variable Number 4.08

Variable Type of operation performed

Variable Name optype

Definition What type of operation was performed for the hip fracture?

Justification To enable the identification of the patient's type of hip fracture operation

Format 2 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 Cannulated screws (e.g. multiple screws)

2 Sliding hip screw

3 Intramedullary nail short 4 Intramedullary nail long

5 Hemiarthroplasty stem cemented 6 Hemiarthroplasty stem uncemented 7 Total hip replacement stem cemented 8 Total hip replacement stem uncemented

97 Other 99 Not known

DD Comments Intramedullary nail includes: Proximal femoral nail, Antegrade femoral nail, Proximal

femoral nail antirotation (PFNA), and Gamma nail.

For cemented versus uncemented procedures, this only includes whether the stem was cemented or not. This does not include whether or not the cup was cemented.

Austin Moore prosthesis to be included in hemiarthroplasty – uncemented.

Sliding hip screws include dynamic hip screws (DHS)

Variable Full weight bear

Variable Name wbear

Definition What is the patient's immediate post-operative weight bearing status?

Justification Ability to monitor variation in practice. Hip Fracture Care Clinical Care Standard

Indicator 5b.

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 0 Unrestricted weight bearing

1 Restricted / non weight bearing

9 Not known

DD CommentsUnrestricted weight bearing refers to a patient who is able to mobilise with full use

of the affected limb to weight bear as pain allows.

Restricted weight bearing refers to a patient where there is a specific instruction that prevents the patient being allowed to fully utilise the leg irrespective of degree of pain. Restricted weight bearing includes terms such as partial weight bear, touch-

weight bear and non-weight bear

Variable First day mobilisation

Variable Name mobil

Definition Was the patient with a hip fracture provided with the opportunity to be mobilised

on day one post hip fracture surgery?

Justification Hip Fracture Care Clinical Care Standard Indicator 5a. Low mobility during

hospitalisation is associated with death, and declining function in activities of daily living at discharge and at one month follow-up, which induces a risk of staying

dependent in these activities (Pedersen et al. 2013).

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 Patient given opportunity to start mobilising day 1 post surgery

1 Patient not given opportunity to start mobilising day 1 post surgery

9 Not known

DD Comments Day 1 post-surgery means the next calendar day following the day of the patient's

primary surgery for hip fracture.

This means the patient was given the opportunity to start mobilising on day 1 post hip fracture surgery. Mobilising may include transferring in/out of bed, stepping, or

walking.

Patients who have been given the opportunity to mobilise but are determined by the clinical team to be too unwell to mobilise are included provided both the opportunity to mobilise and the clinical determination are documented in the

medical record.

Patients who have declined to mobilise are included provided both the opportunity to mobilise and the reason for declining are documented in the medical record.

Pedersen MM, Bodilsen AC, Petersen J, Beyer N, Andersen O, Lawson-Smith L, et al. 2013. Twenty-four-hour mobility during acute hospitalization in older medical patients. The Journals of Gerontology Series A: Biological Sciences and Medical

Sciences 68(3):331-7.

Variable New pressure injuries of the skin

Variable Name Pulcers

Definition Did the patient acquire a new pressure injury (Stage II or above) during their stay in

hospital for the treatment of their hip fracture?

Justification Hip Fracture Care Clinical Care Standard Indicator 5bc Pressure injuries of the skin

are potentially preventable. They can affect a person's level of pain, quality of life,

cost of care, and mortality.

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No

1 Yes

9 Not known

DD Comments A pressure injury is an area of localised damage to the skin and underlying tissue

caused by pressure, shear or friction forces, or a combination of these. Grading for

pressure ulcers consists of 4 levels:

Stage I pressure injury: non-blanchable erythema (intact skin with non-blanchable

redness of a localised area usually over a bony prominence).

Stage II pressure injury: partial thickness skin loss (Partial thickness loss of dermis

presenting as a shallow open ulcer with a red pink wound bed, with slough).

Stage III pressure injury: full thickness skin loss (Subcutaneous fat may be visible but

bone, tendon, or muscle, are not fully exposed).

Stage IV pressure injury: full thickness tissue loss (Full thickness tissue loss with

exposed bone, tendon or muscle).

The pressure injury classification is from the National Pressure Ulcer Advisory Panel (NPUAP) and European Pressure Ulcer Advisory Panel (EPUAP), Prevention and

Treatment of Pressure Ulcers: Clinical Practice Guideline. 2009, Washington DC:

NPUAP

Variable Assessed by geriatric medicine

Variable Name gerimed

Definition Was the patient assessed by geriatric medicine during the acute phase of the

episode of care?

Justification Ability to monitor quality of care. Hip Fracture Care Clinical Care Standard Indicator

3a.

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 0 No

1 Yes

8 No geriatric medicine service available

9 Not known

DD Comments An assessment by geriatric medicine refers to an assessment by a geriatrician or a

medical practitioner (Registrar) working under the supervision of a geriatrician.

The acute phase (IHPA Admitted Hospital Care Types: Guide For Use 2015) is care in

which the primary clinical purpose or treatment goal is to:
• cure illness or provide definitive treatment of injury

perform surgery

• relieve symptoms of illness or injury (excluding palliative care)

• reduce severity of an illness or injury

• protect against exacerbation and/or complication of an illness and/or injury which

could threaten life or normal function

• perform diagnostic or therapeutic procedures

Variable Number 4.14

Variable Geriatric medicine assessment date

Variable Name gdate

Definition Date on which an admitted patient was first assessed by geriatric medicine during

the acute phase of their episode of care

Justification To enable the identification of the date of geriatric assessment. Hip Fracture Care

Clinical Care Standard Indicator 3a.

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments A geriatric assessment is considered to include an assessment by a geriatrician or a

medical practitioner (Registrar) working under the supervision of a geriatrician.

Leave blank if not known

Variable Specialist falls assessment

Variable Name fassess

DefinitionDid the patient undergo a specialist falls assessment? **Justification**Did the patient undergo a specialist falls assessment?

Ability to monitor secondary hip fracture prevention

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 0 No

1 Performed during admission 2 Awaits falls clinic assessment

3 Further intervention not appropriate

8 Not relevant 9 Not known

DD Comments

A specialist falls assessment is undertaken by a multidisciplinary team and includes a systematic assessment by a suitably trained person (i.e. geriatrician or a specialist assessment trained nurse), which must go over the following domains: (i) falls history (noting previous falls); (ii) cause of index fall (including medication review); (iii) risk factors for falling and injury (including fracture) and from this information formulate and document an individualised plan of action to prevent further falls. A specialist falls assessment is not a screening tool.

Example 1: Patient admitted with a hip fracture and managed surgically. During the post-operative period in the acute ward, a specialist falls assessment was commenced with documented assessment of falls risk and medication review. In the medical record there was a documented referral to the specialist falls service to be actioned on discharge from acute care. Option 2 would be selected.

Example 2: Patient admitted with a hip fracture and managed surgically. During the post-operative period in the acute ward, a specialist falls assessment was commenced with documented assessment of falls risk factors, falls history and cause of index fall. There was no other documentation of assessment or referral. Option 0 would be selected.

Variable Bone protection medication at discharge from acute hospital

Variable Name dbonemed1

Definition What bone protection medication was the patient using at discharge from acute

hospital?

Justification Ability to monitor use of bone protection medication. Hip Fracture Care Clinical Care

Standard Indicator 6a.

Format 1 digit numeric

Status Code

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No bone protection medication

1 Yes - Calcium and/or vitamin D only

2 Yes - Bisphosphonates, denosumab, romosozumab or teriparatide (with or

without calcium and/or vitamin D)

9 Not known

DD Comments Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or

one alpha).

Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate,

Zoledronate, Pamidronate.

Variable Delirium assessment

Variable Name delassess

Definition Did the patient have a documented assessment of delirium in the week following

surgery for their hip fracture?

Justification Identifying patients with delirium is the first step in taking action to providing high

quality care. Early diagnosis and prompt treatment offers patients with delirium the

best chance of recovery.

Format 1 digit
Status Non-Core

Coding Source

Coding Frame 1 Not assessed

2 Assessed and not identified3 Assessed and identified

9 Not known

DD Comments

Assessment of delirium requires the use of a validated tool. There are a range of validated diagnostic tools for delirium and they include:

- Confusion Assessment Method (CAM) (Inouye et al. 2014; Shi et al. 2013)
- Confusion Assessment Method (CAM-ICU) (Ely et al. 2001)
- 3D-CAM (Marcantonio et al. 2014).
- The 4AT (Bellelli et al. 2014)

If a person declines assessment record as not assessed.

Delirium is defined as an acute change in mental status that is common among older patients in hospital (Clinical Epidemiology and Health Service Evaluation Unit 2006). It is characterised by a disturbance of consciousness, attention, cognition and perception that develops over a short period of time (usually hours to a few days) (National Institute for Health and Clinical Excellence 2010; Inouye 2014). Patients with delirium may be agitated and restless (hyperactive delirium), quiet and withdrawn (hypoactive delirium), or move between these two subtypes (mixed delirium) (Clinical Epidemiology and Health Service Evaluation Unit 2006; National Institute for Health and Clinical Excellence 2010).

Variable Clinical malnutrition assessment

Variable Name malnutrition

Definition Did the patient undergo clinical assessment of their protein/energy nutrition status

during the acute phase of the episode of care?

Justification Hip fracture patients are at high risk of malnutrition. Malnutrition in these patients

is associated with increased morbidity and mortality, and a decrease in return to

pre-fracture functioning.

Format 1 digit Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 Not done

1 Malnourished2 Not malnourished9 Not known

DD Comments Clinical assessment of a person's nutritional status is encouraged during the acute

phase. Sites should use tools that are validated for such purposes, and are advised to discuss with their Dietitians how best to record the results using this variable's

options.

If the nutritional assessment is performed more than once, please record the first

assessment after admission that uses a validated tool.

Variable First day walking

Variable Name mobil2

DefinitionDid the patient get out of bed and walk on day one post hip fracture surgery? **Justification**Hip Fracture Care Clinical Care Standard Indicator 5a. Low mobility during

hospitalisation is associated with death, and declining function in activities of daily living at discharge and at one month follow-up, which induces a risk of staying

dependent in these activities (Pedersen et al. 2013).

Format 1 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No

1 Yes

9 Not known

DD Comments Day 1 post-surgery means the next calendar day following the day of the patient's

primary surgery for hip fracture. This data item is recording whether the patient

actually stood and stepped or walked by day 1 post-surgery.

Mobilised means the patient managed to stand and step transfer out of bed onto a chair/commode or walk. This does not include only sitting over the edge of the bed

or standing up from the bed without stepping/walking.

Pedersen MM, Bodilsen AC, Petersen J, Beyer N, Andersen O, Lawson-Smith L, et al. 2013. Twenty-four-hour mobility during acute hospitalization in older medical

patients. The Journals of Gerontology Series A: Biological Sciences and Medical

Sciences 68(3):331-7.

Section 5 Discharge

Variable Number 5.01

Variable Discharge date from acute ward

Variable Name wdisch

Definition Date on which the patient was discharged from an acute ward during their episode

of care

Justification To enable the identification of the date of discharge from an acute ward so as to

calculate LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD CommentsThe discharge date refers to the patient physically leaving the acute ward. Record

the date the patient was physically discharged from the acute orthopaedic stay.

Date not known is entered as: 01011900

Variable Number 5.02

Variable Discharge destination from acute orthopaedic episode

Variable Name wdest

Definition What is the discharge (geographical) destination of the patient from the acute/

orthopaedic ward?

Justification To assess patient outcome

Format 2 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 Private residence (including unit in retirement village)

2 Residential aged care facility3 Rehabilitation unit public4 Rehabilitation unit private5 Other hospital / ward / specialty

6 Deceased

7 Short term care in residential care facility (New Zealand only)

97 Other 99 Not known

DD Comments Record the patient's discharge destination at discharge from the acute orthopaedic

stay. If the patient is discharged to live with a relative or in a community group home or boarding house code 'private residence'. Private rehabilitation units will

not be applicable in New Zealand.

Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and accommodation and care for a person on a long-term basis.

in Australia and private hospitals or rest homes in New Zealand.

Short-term care in residential care facility may be relevant if the patient is non-weight bearing, and is used in New Zealand and, to a lesser degree, in Australia. For example, in New Zealand, some District Health Boards have programmes that fund patients who are non-weight bearing to be cared for in a residential care facility. When they are approved by the Orthopaedic Department to fully weight bear they are either admitted to Older Peoples Health for rehabilitation or discharged home.

Variable Discharge from hospital date

Variable Name hdisch

Definition Date on which an admitted patient was discharged from the operating hospital

following their episode of care

Justification To enable the identification of the date of discharge from hospital and calculation of

LOS

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments Date not known is entered as: 01011900

Discharge from hospital date may be the same as discharge from acute ward if patient discharged from hospital system on discharge from acute ward date.

Variable Number 5.04

Variable Length of stay (operating hospital)

Variable Name olos

Definition The length of stay of a patient at the operating hospital, excluding leave days or

days before fracture if occurred in hospital, measured in days

Justification To enable the identification of the length of stay at the operating hospital

Format 3 digit numeric

Status Non-core (created centrally)

Coding Source National Health Data Dictionary, Version 15

Coding Frame NNN

DD Comments Formula: Length of Stay (LOS) = Separation date - Admission date - Total leave days.

The calculation is inclusive of admission and separation dates. LOS will be calculated

automatically from the operating hospital separation and admission dates.

If the hip fracture occurred as an in-patient then the length of stay should be from

time hip fracture was diagnosed.

Variable Length of stay (health system)

Variable Name TLOS

Definition The length of stay of a patient from admission/diagnosis of a hip fracture to final

date of discharge from an inpatient facility (public or private), excluding leave days,

measured in days

Justification To enable the identification of the total length of stay in the health system

Format 4 digit, unit of measure (day)

Status Non-core

Coding Source National Health Data Dictionary, Version 15

Coding Frame NNNN

DD Comments Formula: Length of stay (LOS) = Separation date - Admission date - Total leave days.

The calculation is inclusive of admission and separation dates.

LOS will be calculated automatically from the ED arrival date of the transferring hospital (or the ED arrival date of the operating hospital, if no transfer occurred) and the discharge from hospital date. If the final date of discharge from the hospital

system is known, this date should be used.

It should be noted that the total length of stay in the hospital system will be difficult

to calculate in some jurisdictions, due to differences in treatment settings for

rehabilitation-based care.

Variable Discharge place of residence

Variable Name dresidence

Definition What is the usual place of residence of the person following discharge from the

whole hospital system?

Justification Type of accommodation before and after admission are collected to compare where

the patient has come from (what was their usual accommodation) and where they are going to (what will become their usual accommodation). Comparison of accommodation pre and post admission is an indicator of patient outcome.

1 digit numeric

Status Core

Format

Coding Source Adapted from Australasian Rehabilitation Outcomes Centre Inpatient Dataset,

Version 3.0; NSW SNAP Data Collection, Version 4.0

Coding Frame 1 Private residence (including unit in retirement village)

2 Residential aged care facility

3 Deceased7 Other9 Not known

DD Comments Record the patient's accommodation type at discharge from the whole hospital

system.

If the patient lives with a relative or in a community group home or boarding house

code 'private residence'.

Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services

in Australia and private hospitals or rest homes in New Zealand.

Section 7 120 day follow-up*

*120-day follow up is undertaken by the operating hospital

Variable Number 7.01

Variable 120 day follow-up date

Variable Name fdate2

Definition Date on which the 120 day follow-up was completed post the initial hip fracture

surgery

Justification To monitor patient outcomes post-surgery

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments Date not known is entered as: 01011900

Variable Number 7.02

Variable Survival at 120 days post-surgery

Variable Name fsurvive2

DefinitionIs the patient alive at 120 days post-surgery**Justification**To monitor patient outcomes post-surgery

Format 1 digit numeric

Status Core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

DD Comments

Variable Number 7.03

Variable Date health system discharge at 120 day follow-up

Variable Name date120

Definition What date was the patient discharged from the hospital system?

Justification To enable the identification of the total length of stay in the health system

Format 8 digit date, date in DDMMYYYY

Status Core

Coding Source National Health Data Dictionary, Version 15

Coding Frame DD/MM/YYYY

DD Comments

Variable Place of residence at 120 day follow-up

Variable Name fresidence2

Definition What is the place of residence of the person at 120 days post-surgery?

Justification To monitor patient outcomes post-surgery. Hip Fracture Care Clinical Care Standard

Indicator 7b.

Format 2 digit numeric

Status Core

Coding Source Adapted from the Australasian Rehabilitation Outcomes Centre Inpatient Dataset,

Version 3.0; NSW SNAP Data Collection, Version 4.0

Coding Frame 1 Private residence (including unit in retirement village)

2 Residential aged care facility3 Rehabilitation unit public4 Rehabilitation unit private5 Other hospital / ward / specialty

6 Deceased

7 Short term care in residential care facility (New Zealand only)

97 Other 99 Not known

DD Comments Record the patient's discharge destination at 120 days post-surgery. If the patient is

discharged to live with a relative or in a community group home or boarding house code 'private residence'. Private rehabilitation units will not be applicable in New

Zealand.

Residential aged care facility refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-

purpose services in Australia and private hospitals or rest homes in New Zealand.

Short-term care in residential care facility may be relevant if the patient is non-weight bearing and is used in New Zealand and to a lesser degree in Australia. For example, in New Zealand, some District Health Boards have programmes that fund patients who are non-weight bearing to be cared for in a residential care facility. When they are approved by the Orthopaedic Department to fully weight bear they are either admitted to Older Peoples Health for rehabilitation or discharged home.

Variable Post-admission walking ability at 120 day follow-up

Variable Name fwalk2

Definition What was the patient's walking ability at 120 days post-surgery?

Justification To monitor patient mobility status post-discharge. Hip Fracture Care Clinical Care

Standard Indicator 5d.

Format 2 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 1 Usually walks without walking aids

2 Usually walks with either a stick or crutch3 Usually walks with two aids or frame4 Usually uses a wheelchair / bed bound

8 Not relevant 9 Not known

DD CommentsUsually walks with two aids or frame includes with or without assistance of a person

If a person has different levels of mobility on different surfaces then record the level of most assistance. For example, inside their residence a person usually walks with no walking aid but when outside the residence the person usually walks with a

frame, then the level of mobility recorded is option 3.

Variable Bone protection medication at 120 day follow-up

Variable Name fbonemed2

Definition What bone protection medication was the patient using at 120 days post-surgery?

Justification Ability to monitor use of bone protection medication

Format 1 digit numeric

Status Code

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No bone protection medication

1 Yes - Calcium and/or vitamin D only

2 Yes - Bisphosphonates, denosumab, romosozumab or teriparatide (with or without

calcium and/or vitamin D)

9 Not known

DD Comments Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or

one alpha).

Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate,

Zoledronate, Pamidronate.

Variable Number 7.08

Variable Re-operation within 120 day follow-up

Variable Name fop2

Definition What kind of re-operation has been required (if any) for the patient within 120 days

post-surgery?

Justification To monitor patient outcomes post-surgery

Format 2 digit numeric

Status Core

Coding Source Adapted from the UK National Hip Fracture Database

Coding Frame 0 No reoperation

1 Reduction of dislocated prosthesis

2 Washout or debridement

3 Implant removal

4 Revision of internal fixation 5 Conversion to hemiarthroplasty 6 Conversion to total hip replacement

7 Excision arthroplasty 9 Revision arthroplasty

99 Not known

DD Comments Option 2 washout and debridement includes liner change. Note: record the most

significant procedure only.

Variable Preliminary date of death

Variable Name predod

Definition What was the date of death of the hip fracture patient?

Justification To monitor patient outcomes and enable reporting of mortality after hip fracture

Hip Fracture Care Clinical Care Standard Indicator 8b.

Format 8 digit, date in DDMMYYYY

Status Optional, non-core

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 646025).

Preliminary Australian date of death obtained from hospital records and/or during 120 day

follow-up.

Coding Frame DD/MM/YYYY

DD Comments

Date of death may be collected either at discharge or during 120-day follow-up.

New Zealand date of death may be obtained from the New Zealand Ministry of Health.

Variable Number 7.10

Variable Final date of death

Variable Name findod

Definition What was the date of death of the hip fracture patient?

Justification To monitor patient outcomes and enable reporting of mortality after hip fracture

Hip Fracture Care Clinical Care Standard Indicator 8b.

Format 8 digit, date in DDMMYYYY

Status Non-core (created centrally)

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 646025).

Final Australian date of death obtained from the National Death Index.

New Zealand date of death obtained from the New Zealand Ministry of Health.

Coding Frame DD/MM/YYYY

DD Comments

Final Australian date of death will be obtained from the National Death Index and final New

Zealand date of death will be obtained from the New Zealand Ministry of Health.

Variable Underlying cause of death

Variable Name undcod

Definition What was the underlying cause of death of the hip fracture patient?

Justification To enable identification of the underlying cause of death of the hip fracture patient

Format ANN {.N [N]}

Status Non-core (created centrally)

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 307862).

Australian underlying cause of death obtained from the National Death Index.

New Zealand underlying cause of death obtained from the New Zealand Ministry of Health.

Coding Frame ICD-10

DD Comments The disease or injury which initiated the train of morbid events leading directly to a person's

death or the circumstances of the incident or violence which produced the fatal injury.

Variable Number 7.12

Variable Other causes of death

Variable Name othcod

Definition What was the underlying cause of death of the hip fracture patient?

Justification To enable identification of the underlying cause of death of the hip fracture patient

Format ANN {.N [N]}

Status Non-core (created centrally)

Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 307862).

Australian other cause(s) of death obtained from the National Death Index.

New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.

Coding Frame ICD-10

DD Comments The disease or injury which initiated the train of morbid events leading directly to a person's

death or the circumstances of the incident or violence which produced the fatal injury.

EQ5D5L questionnaire

UK (English) © 2009 EuroQol Group EQ-5D $^{\mathrm{m}}$ is a trade mark of the EuroQol Group


Under each heading, please tick the ONE box that best describes your health TODAY.

MOBILITY		
I have no problems in walking about		?
have slight problems in walking about		?
have moderate problems in walking about	?	
have severe problems in walking about		?
l am unable to walk about		?
SELF-CARE		
have no problems washing or dressing myself		?
have slight problems washing or dressing myself	?	
have moderate problems washing or dressing myse	elf	?
have severe problems washing or dressing myself	?	
am unable to wash or dress myself		?
USUAL ACTIVITIES to a work study bousework fam	silv or loid	ura activitias)
USUAL ACTIVITIES (e.g. work, study, housework, fan I have no problems doing my usual activities	illy of leis	are activities)
have slight problems doing my usual activities		[?]
have moderate problems doing my usual activities	ы	ī.
have severe problems doing my usual activities	E)	[?]
am unable to do my usual activities		?
and unable to do my usual activities		E
PAIN / DISCOMFORT		
have no pain or discomfort		?
have slight pain or discomfort		?
have moderate pain or discomfort		?
have severe pain or discomfort	?	
have extreme pain or discomfort	?	
ANXIETY / DEPRESSION		
am not anxious or depressed		?
am slightly anxious or depressed	?	ш
am moderately anxious or depressed	ш	?
am severely anxious or depressed		?
am extremely anxious or depressed		?
an extremely anxious or depressed		ш

The best health you can imagine

- We would like to know how good or bad your health is TODAY.
- . This scale is numbered from 0 to 100.
- 100 means the <u>best</u> health you can imagine.
 0 means the <u>worst</u> health you can imagine.
- Mark an X on the scale to indicate how your health is TODAY.
- Now, please write the number you marked on the scale in the box below.

YOUR HEALTH TODAY =



ANZHFR Facility Level Audit

Hospitals are identified using the variable 2.01: Establishment identifier of operating hospital

Section 8 Hospital information		
Variable # Variable	8.01 Major trauma centre	
	·	
Variable Name	- -	
Definition	Is the hospital a designated major trauma centre?	
Justification	To identify the Level 1 trauma centres	
Format	Numerical, N	
Status	non core	
Coding Source		
Coding Frame	0 No 1 Yes	
	9 Not known	
FLA Comment	Investigators can determine this using the Australasian trauma verification program manual. The manual is available at: https://www.surgeons.org/media/21043200/march-2016-trauma-verification-manual.pdf	
Variable #	8.02	
Variable	Hip fractures	
Variable Name	est_numb_hipfrac	
Definition	Estimated number of hip fractures in the calendar year just ended January to December inclusive	
Justification	To estimate the number of hip fractures being treated at the hospital	
Format	Numerical, NNNN	
Status	core	
Coding Source		
Coding Frame	1 0-50	
	2 51-100	
	3 101-150 4 151-200	
	5 201-300	
	6 301-400	
	7 401+	
	9 Not known	
FLA Comment	s Record the estimated number of fractures treated annually.	

Section 9 Model of care

Variable # 9.01

Variable Orthogeriatric service

Variable Name ogs

Definition Was there a formal orthogeriatric service in place?

Justification To determine if there was an orthogeriatric service available for hip fracture patients at the

hospital

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments Orthogeriatric care involves a shared care arrangement of hip fracture patients between the

specialties of orthopaedics and geriatric medicine. The geriatrician is involved in the preoperative optimisation of the patient in preparation for surgery and then takes a lead in the post-operative medical care and coordinates the discharge planning process. Implicit in this role are many of the aspects of basic care including nutrition, hydration, pressure care, bowel and bladder management, and monitoring of cognition (ANZHFR Guideline 2014, p.68).

54

Variable # 9.02

Variable Model of care

Variable Name m

Definition Select the model of care that best describes the service provided for care of older hip fracture

patients in your hospital.

Justification To determine the model of care used to treat hip fracture patients. Hip Fracture Care Clinical

Care Standard Indicator 3a.

Format Numerical, N

Status core

Coding Source

Coding Frame 1 Orthopaedics and Geriatric Medicine shared care

2 Orthogeriatric Liaison Service where Geriatric Medicine provides daily review during working

week

3 Medical Liaison Service where Physician or GP provide daily review during working week

4 Orthogeriatric Liaison Service where Geriatric Medicine provides intermittent review once or more per week

5 Medical Liaison Service where Physician or GP provides intermittent review once or more per week

6 A geriatric service provided on referral 7 A Medical Service provided on referral

8 No formal service

9 Other 99 Not known

FLA Comments

Documented local arrangements for the management of hip fracture patients according to an orthogeriatric (or alternative physician or medical practitioner) model of care. The documentation should be an agreement showing acceptance of a "shared care" model for all hip fracture patients, and signed by the heads of both Geriatric Medicine and Orthopaedic Surgery.

The key features of an orthogeriatric model of care are:

- regular medical assessment including medication review;
- managing patient comorbidities;
- optimisation for surgery;
- early identification of each patient's goals and care co-ordination. If appropriate and clinically indicated, provision of multidisciplinary rehabilitation aimed at increasing mobility and independence, and to facilitate a return to pre-fracture residence and support long-term wellbeing;
- early identification of most appropriate service to deliver rehabilitation, if indicated;
- ongoing orthogeriatric and multidisciplinary review including reassessment of cognition after surgery, and discharge planning liaison with primary care, including falls prevention and secondary fracture prevention.

Section 10 Protocols and processes

Variable # 10.01

Variable Imaging protocol

Variable Name ct_mri

Definition For a suspected hip fracture, does your hospital have a protocol or pathway for access to CT /

MRI for inconclusive plain imaging?

Justification To determine if the hospital has a protocol for the imaging of patients suspected of having a

hip fracture

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments CT is Computed Tomography

MRI is Magnetic Resonance Imaging

Offer magnetic resonance imaging (MRI) if hip fracture is suspected despite negative anteroposterior pelvis and lateral hip X-rays. If MRI is not available within 24 hours or is

contraindicated, consider computed tomography (CT).

Variable # 10.02

Variable Hip fracture pathway

Variable Name hipfrac path

Definition The hospital has a hip fracture pathway that is used for the management of patients admitted

with a hip fracture.

Justification To determine if the hospital has a hip fracture pathway. Hip Fracture Care Clinical Care

Standard Indicator 1a.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes - ED only

2 Yes - whole acute journey

9 Not known

FLA Comments Evidence of local arrangements for the management of patients with hip fracture in the

emergency department. Documented local arrangements for the management of patients with hip fracture in the emergency department that address timely assessment and management of the patient's medical conditions, including but not limited to: diagnostic imaging; pain control; cognitive assessment. The documentation may be in the form of local

protocols and/or a clinical pathway.

Variable Venous thromboembolism protocol

Variable Name vte

Definition Does your hospital have a VTE protocol?

Justification To determine if the hospital has a VTE protocol for hip fracture patients

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments VTE refers to venous thrombo-embolism

Variable # 10.04

Variable Pain protocol
Variable Name pain_path_def

Definition Does your hospital have a protocol or pathway for the management of pain in hip fracture

patients?

Justification To determine if the hospital has a pain protocol for hip fracture patients. Hip Fracture Care

Clinical Care Standard Indicator 2a.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes ED only

2 Yes whole acute journey

9 Not Known

FLA Comments Documented local arrangements include a written clinical protocol to ensure patients with a

hip fracture receive prompt and effective pain management. The protocol should take into account the hierarchy of pain management medicine for managing pain associated with hip

fracture and aim to minimise the use of opioid medicine.

Pain should be assessed immediately upon presentation at hospital and within 30 minutes of administering initial analgesia and hourly until settled on the ward and regularly as part of

routine nursing observations throughout admission.

Protocols should include the use of a standardised pain assessment system, which specifically addresses the assessment of pain for patients with cognitive impairment and those unable to communicate pain, particularly with regard to minimising the use of opioid medicine in this

group.

Variable Planned theatre list

Variable Name oplist_planned

Definition Does your hospital have a planned emergency list / planned orthopaedic trauma list for hip

fracture patients?

Justification To determine if the hospital has access to an appropriately skilled operating team for patients

admitted with a hip fracture.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments A planned emergency list or planned orthopaedic trauma list provides access to an

appropriately skilled team to undertake the surgical procedure.

Variable # 10.06

Variable Anaesthesia

Variable Name anaes_choice

Justification To determine if the hospital routinely offers a choice of anaesthesia for hip fracture patients

Format Numerical, N

Status core

Coding Source

Coding Frame 0 Never

1 Rarely2 Frequently3 Always9 Not known

FLA Comments Patients should be involved in the decision as of the approach to anaesthesia taken. They

should be made aware of the potential risks and benefits of both general and regional

anaesthesia so as to be able to make an informed decision about their care.

Variable Nerve block for pain pre-surgery

Variable Name nvblock_preop

urgery?

Justification To determine if the hospital offers hip fracture patients local nerve blocks as part of pain

management pre-surgery

Format Numerical, N

Status core

Coding Source

Coding Frame 0 Never

1 Rarely 2 Frequently 3 Always 9 Not known

FLA Comments Nerve blocks offer an alternative to systemic analgesia and have the potential to reduce the

dose requirements of potent systemic analgesic agents, which may reduce unwanted side

effects such as sedation, respiratory complications and delirium.

Variable # 10.08

Variable Nerve block for pain post-surgery

Variable Name nvblock_postop

Definition Are local nerve blocks used at the time of surgery to help with postoperative pain?

Justification To determine if the hospital offers hip fracture patients local nerve blocks as part of pain

management post-surgery

Format Numerical, N

Status core

Coding Source

Coding Frame 0 Never

1 Rarely2 Frequently3 Always9 Not known

FLA Comments Nerve blocks offer an alternative to systemic analgesia and have the potential to reduce the

dose requirements of potent systemic analgesic agents, which may reduce unwanted side

effects such as sedation, respiratory complications and delirium.

Variable Therapy access
Variable Name therapy_we_def

Definition Does your hospital offer hip fracture patients routine access to therapy services at weekends?

Justification To determine if the hospital offers hip fracture patients therapy services at weekends

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes - Physiotherapy only

2 Yes – other 9 Not known

FLA Comments Early mobilisation is also associated with short term gains related to a reduction in

postoperative complications. Unless medically or surgically contraindicated, mobilisation should start the day after surgery. Patients should be offered an opportunity to mobilise at

least once a day with regular physiotherapy review ensured.

Section 11 Beyond the Acute Hospital Stay

Variable # 11.01

Variable Information on treatment and care

Variable Name hipfrac_written

Definition Does your hospital routinely provide patients and/or family and carers with written

information about treatment and care for a hip fracture?

Justification To determine if the hospital routinely provides hip fracture patients and/or their family/carers

with written information about their hip fracture treatment and care

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments

Variable # 11.02

Variable Inpatient rehabilitation

Variable Name inpt_rehab

Definition Access to in-patient rehabilitation

Justification To determine if the hospital provides on- or off-site hip fracture rehabilitation for patients

unable to meet the criteria for early supported discharge

Format Numerical, N

Status core

Coding Source

Coding Frame 1 Onsite

2 Offsite 3 Both

4 No inpatient rehabilitation available

9 Not known

FLA Comments Consider in-patient rehabilitation for those in whom further improvement with a structured

multidisciplinary programme is anticipated.

Variable Home-based rehabilitation

Variable Name homebased_serv

Definition Does your hospital have access to an early supported home-based rehabilitation service (not

the same as the Commonwealth funded transitional aged care program or community

services)?

Justification To determine if the hospital has access to early supported home-based hip fracture

rehabilitation for patients recovering from a hip fracture.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments Early supported discharge should be considered provided the patient is medically stable and

has the mental ability to participate in continued rehabilitation and is able to transfer and mobilise short distances and has not yet achieved their full rehabilitation potential, as

discussed with the patient, carer and family.

Variable Injury prevention

Variable Name prevention_written

Definition Does your service provide individualised <u>written</u> information to patients on discharge that

includes recommendations for future falls and fracture prevention? (not the same as a copy of

a discharge summary)

Justification To determine if the hospital provides written information to patients on discharge regarding

fall and fracture-related injury prevention. Hip Fracture Care Clinical Care Standard Indicator

7a.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments Evidence of local arrangements for the development of an individualised care plan for hip

fracture patients prior to the patient's separation from hospital. Documented local arrangements for patients with a hip fracture to have an individualised care plan developed prior to the patients separation from hospital, and provisions to make this available to them (and/or their carer), and to their general practitioner and other ongoing clinical care provider

within 48 hours of the patient leaving the hospital.

The plan should describe the care received by the patient during their hospital stay and ongoing care and goals of care. The plan must include a summary of any changes to medicines, any new medicines, and equipment and contact details for rehabilitation services they may require. It should also describe mobilisation activities, wound care and function post-surgery,

and include information and recommendations for secondary fracture prevention.

Variable # 11.05
Variable Falls clinic

Variable Name falls_clinic

Definition Does your service have access to a Falls Clinic (Public)

Justification To determine if the hospital has access to a Falls clinic for the prevention of future falls

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

Variable Osteoporosis clinic

Variable Name op_clinic

Definition Does your service have access to an Osteoporosis Clinic (Public)

Justification To determine if the hospital has access to an osteoporosis clinic for the management of bone

health

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments

Variable # 11.07

Variable Falls and bone health clinic

Variable Name

falls_bone_clinic_comb

Definition Does your service have access to a combined Falls and Bone Health Clinic (Public)

Justification To determine if the hospital has access to a Falls and Bone Health clinic for the management

and prevention of future injury.

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments

Variable # 11.08

Variable Orthopaedic clinic

Variable Name ortho_clinic

Definition Does your service have access to an Orthopaedic Clinic (Public)

Justification To determine if the hospital has access to an Orthopaedic clinic

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

Variable Fracture liaison service
Variable Name frac_liaison_serv_def

Definition Do you have a Fracture Liaison Service, whereby there is systematic identification of fracture

patients by a fracture liaison nurse, with a view to onward referrals and management of

osteoporosis?

Justification To determine if the hospital has access to a fracture liaison service

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 Yes – hip fracture patients only

2 Yes – all fracture patients (including hip)

9 Not known

FLA Comments A Fracture Liaison Service may employ health care professionals who are not nurses, such as

physiotherapists, and who are called Fracture Liaison Coordinators.

Section 12 Other aspects of care

Variable # 12.01

Variable Hip fracture data

Variable Name data_collect

Definition Does your hospital routinely collect hip fracture data?

Justification To determine if the hospital routinely collects hip fracture data to enable review of service

provision and outcomes

Format Numerical, N

Status core

Coding Source

Coding Frame 0 No

1 ANZ Hip Fracture Registry

2 Local System 9 Not known

FLA Comments

Variable # 12.02

Variable Service provision plans

Variable Name serv_alt_12mths

Definition Do you have any plans to alter any of your service provision for hip fracture patients over the

next 12 months - if so please give details?

Justification To determine if the hospital will alter any service provision for hip fracture patients

Format Numerical, N Status non core

Coding Source

Coding Frame 0 No 1 Yes

9 Not known

FLA Comments

Variable # 12.03

Variable Service provision plan details

Variable Name serv_alt_detail

Definition Type of service provision plans

Justification To determine the type of service provision changes that are to be made

Format Text Status non core

Coding Source Coding Frame FLA Comments

Variable Service provision barriers

Variable Name serv_alt_barriers

Justification To determine if there are any perceived barriers to service provision changes

Format Numerical, N Status non core

Coding Source

Coding Frame 0 No

1 Yes

9 Not known

FLA Comments

Variable # 12.05

Variable Service provision barrier details

Variable Name serv_barriers_detail

Definition Type of barriers to proposed service redesign

Justification To determine the type of perceived barriers to service provision changes

Format Text Status non core

Coding Source Coding Frame FLA Comments

Variable # 12.06

Variable Hip fracture data collector

Variable Name Data_collector

Definition Who currently collects the data?

Justification To examine the resources required for participation in the ANZHFR

Format Text Status non core

Coding Source

Coding Frame 1 Orthopaedic surgeon

2 Geriatrician

3 Fracture Liaison Coordinator

4 Dedicated Neck of femur nurse/coordinator

5 Specialist orthopaedic nurse 6 Research nurse / coordinator

7 Allied Health

8 Other (Please specify)

9 Not applicable

Variable Hip fracture data entry

Variable Name Data_entry

Definition Does the same person who collects the data enter it into the database/Registry?

Justification To examine the resources required for participation in the ANZHFR

Format Text Status non core

Coding Source

Coding Frame 0 No

1 Yes

9 Not applicable

FLA Comments

-____

_

Variable # 12.08

Variable Time taken data collection and entry

Variable Name Time_core

Definition How long do you estimate data collection and entry take per patient, excluding 120-day

follow-up?

Justification To examine the resources required for participation in the ANZHFR

Format Text Status non core

Coding Source

Coding Frame 1 Less than 30 minutes per patient

2 30 minutes per patient3 45 minutes per patient4 60 minutes per patient5 75 minutes per patient

6 > 75 minutes per patient (Please specify)

9 Not applicable

FLA Comments

Variable # 12.09

Variable Time taken 120-days

Variable Name Time_120

Definition If you undertake 120-day follow-up, how long do you estimate data collection and entry take

per patient?

Justification To examine the resources required for participation in the ANZHFR

Format Text Status non core

Coding Source

Coding Frame 1 Less than 30 minutes per patient

2 30-60 minutes per patient3 61-90 minutes per patient

4 >90 minutes per patient (Please specify)

9 Not applicable

Variable # 12.10

Variable Sprint audits

Variable Name sprints

Definition Would your hospital be interested in participating in future ANZHFR Sprint Audits?

Justification To explore support for voluntary participation in Sprint Audits

Format Text Status non core

Coding Source

Coding Frame 0 No

1 Yes 2 Maybe

FLA Comments The ANZHFR introduced Sprint Audits in 2021. A Sprint Audit involves additional

questions/variables that are temporarily added to the routine registry data collection. Sprint

Audits allow us to get a 'close up' view of specific aspects of hip fracture care.

.

Variable # 12.11

Variable Impact of COVID Variable Name covid_changes

Definition Were there any changes in the way older patients with a hip fracture were cared for in your

hospital during 2020 due to the impact of COVID-19 on your service?

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 No

1 Yes

FLA Comments

Variable Impact of COVID

_

Variable # 12.12

Variable Impact of COVID: Local arrangements for the management of patients with hip fracture in

the ED

Variable Name care_presentation_a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact2 No discernible impact

Variable Impact of COVID: Assessment of preoperative cognitive status

Variable Name care_presentation_b

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact2 No discernible impact

FLA Comments

Variable # 12.14

Variable Impact of COVID: Pain assessment

Variable Name pain_mx_a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.15

Variable Impact of COVID: Pain management

Variable Name pain_mx_b

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

Variable Impact of COVID: Orthorgeriatric model of care

Variable Name og_moc_a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text
Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.17

Variable Impact of COVID: Time to surgery

Variable Name tts a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.18

Variable Impact of COVID: Day one mobilisation

Variable Name mob wb a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.19

Variable Impact of COVID: Unrestricted weightbearing

Variable Name mob_wb_b

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact2 No discernible impact

FLA Comments

Variable # 12.20

Variable Impact of COVID: Pressure injuries

Variable Name mob_wb_c

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.21

Variable Impact of COVID: Return to pre-fracture mobility

Variable Name mob_wb_d

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text

Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact2 No discernible impact

FLA Comments

Variable # 12.22

Variable Impact of COVID: Bone protection medication at discharge

Variable Name min_risk_frac_a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact2 No discernible impact

FLA Comments

Variable # 12.23

Variable Impact of COVID: Local arrangements for the development of an individualised care plan for

hip fracture patients prior to the patient's separation

Variable Name trans_hosp_a

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact

FLA Comments

Variable # 12.24

Variable Impact of COVID: Proportion of patients returning to private residence within 120 days post

separation from hospital

Variable Name trans_hosp_b

Definition Considering each of the Hip Fracture Care Clinical Care Indicators, please select the option that

best describes how you feel that particular Indicator was impacted by changes due to COVID-

19.

Justification To determine if there were any changes in hip fracture care due to the COVID-19 pandemic

Format Text Status non core

Coding Source

Coding Frame 0 Negative impact

1 Positive impact

2 No discernible impact