



ANZHF

Australian & New Zealand Hip Fracture Registry

Data Dictionary
Version 17

October 2024

Australian and New Zealand Hip Fracture Registry

Background: A minimum data set (MDS) was created for the Australian and New Zealand (ANZ) Hip Fracture Registry Steering Group. The purpose of the minimum dataset and registry is to support consistent, local collection of data across Australia and New Zealand to enable facilities to compare practice and outcomes for hip fracture care against national clinical guidelines and standards of care. The MDS has been developed to capture information relevant to ANZ Hip Fracture Guidelines and national Hip Fracture Care Clinical Care Standard and is comparable to the United Kingdom (UK) national hip fracture registry and other registries emerging across the world.

Purpose: The ultimate goal of the ANZ Hip Fracture Registry is to use data to improve performance and maximise outcomes for older people who fracture their hip by reducing mortality and morbidity, reducing rates of institutionalisation, maximising functional independence and preventing future fractures by monitoring secondary prevention interventions.

MDS development: The MDS has been reviewed by the ANZ Hip Fracture Registry Steering Group, which consists of representatives of key professional and consumer bodies from Australia and New Zealand: Australian and New Zealand Society for Geriatric Medicine (ANZSGM); Australian Orthopaedic Association (AOA); Australian and New Zealand College of Anaesthetists (ANZCA); Australasian College of Emergency Medicine (ACEM); New Zealand Orthopaedic Association (NZOA); Royal Australasian College of Surgeons (RACS); Royal Australasian College of Physicians (RACP); Australian and New Zealand Orthopaedic Nurses Association (ANZONA); Australasian Faculty of Rehabilitation Medicine (AFRM); Australian Physiotherapy Association (APA); Osteoporosis Australia (OA); and Osteoporosis New Zealand (ONZ). This version of the ANZHFR Data Dictionary includes data variables for both the Patient Level Audit (the Registry) and the Facility Level Audit (annual snapshot of hospital level processes and protocols).

The data variables collected in the MDS (Patient Level) are from six (6) key components of care and include: (1) Patient information; (2) Admission; (3) Assessment; (4) Treatment; (5) Discharge; and (7) 120 day follow-up. The data variables collected in the MDS (Facility Level) cover: (1) Hospital Information; (2) Model of Care; (3) Protocols and processes; (4) Beyond the acute hospital stay; (5) Other aspects of care.

Core and non-core data items

Core variables are those variables collected by all using the minimum dataset e.g. date and time of admission, or type of hip fracture, and will be uploaded to the ANZ Hip Fracture Registry (ANZHFR). A number of these items will be considered mandatory for the purposes of forming a meaningful registry. Non-core items are collected at a local level and are held either locally or on the central server, or are generated automatically at a central level using data uploaded.

Review: The MDS will be reviewed annually by the ANZHFR Steering Group. It is anticipated that any new item to be added must be presented with a clear case for the benefits of adding it. Equally the Committee will be charged with removing redundant items which are felt not to add value either at a facility or central level.

Patient Inclusion: A person aged 50 years or older, who has been admitted to a participating hospital with an acutely fractured hip from a minimal or low trauma injury, and who undergoes either surgical or non-surgical management of the hip fracture.

Version history:

Version	Description of Change	Author	Date Changed	Status
1.0	Draft	Rebecca Mitchell	July 2012	Rough draft
2.0	Draft	Rebecca Mitchell	6 Dec 2012	Draft
3.0	Draft	Rebecca Mitchell	8 Jan 2013	Draft
4.0	Draft	Rebecca Mitchell	3 May 2013	Draft
5.0	Draft	Rebecca Mitchell	31 May 2013	Draft
6.0	Penultimate	Rebecca Mitchell	21 June 2013	Draft
7.0	Final	Rebecca Mitchell	5 July 2013	Final
8.0	Revised to ensure data capture consistent with Guideline recommendations	Jacqui Close	4 Dec 2013	Final
8.1	Revised to ensure data capture consistent with Guideline recommendations and the	Jacqui Close	11 Dec 2013	Final

	requirement to capture identifying data for follow up and data linkage			
9.0	Review by the Steering Group against the 2014 ANZ Guidelines for Hip Fracture Care and the 2016 ACSQHC Hip Fracture Care Clinical Care Standard and Indicators; incorporation of definitions for the Facility Level Audit variables	Elizabeth Armstrong	August 2016	Draft
9.1	Revision with Steering Group and Data Committee feedback	Elizabeth Armstrong	September 2016	Final Draft
10.0	Review by the Steering Group to incorporate feedback from participating sites and ensure data dictionary continues to be fit for purpose	Steering Group	August 2017	Draft
10.1	Revision with Data Committee feedback	Data Committee	October 2017	Draft
10.2	Revision with Data Committee feedback	Data Committee	October 2017	Final Draft
11	Annual Steering Group review to ensure data dictionary continues to be fit for purpose	Data Committee	October 2018	Final
12	Annual Steering Group review to ensure data dictionary continues to be fit for purpose	Data Committee	October 2019	Final Draft
13	Annual review of the dataset	Data Committee	October 2020	Final
14	Annual review of the dataset	Data Committee	October 2021	Final
15	Annual review of the dataset	Data Committee	October 2022	Final
16	Annual review of the dataset	Data Committee	November 2023	Final
17	Annual review of the dataset	Data Committee	October 2024	Final

List of Data Variables

ANZHFR Patient Level Audit	6
Section 1 Patient information	6
Unique identifier.....	6
Australian and New Zealand jurisdiction.....	6
First name of patient.....	7
Surname of patient.....	7
Hospital MRN / URN / event number.....	7
Contact telephone number for patient.....	8
Date of birth.....	8
Age derived.....	8
Sex of person.....	9
Australian Indigenous status	9
NZ ethnic status.....	10
Patient's postcode.....	111
Medicare number (Australia) / National Health Index (New Zealand)	111
Patient type.....	12
Usual place of residence	13
Statistical linkage key 581	13
Patient email.....	13
Individual Health Identifier.....	14
Section 2 Admission	15
Establishment identifier of operating hospital.....	15
Admission via ED of operating hospital.....	15
Transfer hospital.....	16
ED / hospital arrival date (transfer hospital).....	16
ED arrival time (transfer hospital)	17
ED / other ward arrival date (operating hospital)	17
ED / other ward arrival time (operating hospital)	18
ED departure date (operating hospital)	18
ED departure time (operating hospital)	18
In-patient fracture date.....	19
In-patient fracture time.....	19
Pain management.....	20
Nerve block before transfer	Error! Bookmark not defined.
Section 3 Assessment	22
Pre-admission walking ability.....	22
Pre-operative cognitive assessment	22
Pre-admission cognitive status	23
Bone protection medication at admission	23
Pre-operative medical assessment	24
Side of hip fracture.....	24
Atypical fracture.....	24
Type of fracture.....	25
Surgical repair.....	25
ASA grade.....	26
Clinical Frailty Scale	27
Delirium assessment prior to surgery	27
Section 4 Treatment	29
Date of surgery for hip fracture	29
Time of surgery for hip fracture	29
Surgery delay.....	30
Surgery delay other text.....	30
Type of anaesthesia.....	31
Analgesia - nerve block	31
Consultant surgeon present.....	32
Type of operation performed.....	32
Full weight bear.....	33

New pressure injuries of the skin	33
Assessed by geriatric medicine	34
Geriatric medicine assessment date	344
Bone protection medication at discharge from acute hospital.....	36
Postoperative delirium assessment	37
Clinical malnutrition assessment.....	38
First day walking.....	38
Oral nutritional supplements.....	39
Section 5 Discharge	400
Discharge date from acute ward.....	40
Discharge destination from acute ward	40
Discharge from hospital date	41
Length of stay (acute ward).....	41
Length of stay (hospital system)	42
Discharge place of residence.....	43
Section 7 120 day follow-up*	44
120 day follow-up date	44
Survival at 120 days post-surgery	44
Date hospital system discharge at 120 day follow-up	44
Place of residence at 120 day follow-up	45
Post-admission walking ability at 120 day follow-up	46
Bone protection medication at 120 day follow-up	46
Re-operation within 120 day follow-up	Error! Bookmark not defined.
Preliminary date of death	47
Final date of death.....	48
Underlying cause of death	48
Other causes of death	49
EQ5D5LQuestionnaire.....	52
Section 13 52-week follow-up*	55
Follow-up at 52 weeks	44
52 week follow-up date.....	55
52 week residence.....	55
52 week mobility.....	56
52 week medication	56
Reason for no medication at 52 weeks	57
Further falls.....	57
Strength and balance	58
Further fractures.....	58

ANZHFR Patient Level Audit

Section 1	Patient information
Variable Number	1.01
Variable	Unique identifier
Variable Name	ID
Definition	A consecutive number allocated to each record of a hip fracture
Justification	To allow for the identification of records
Format	10 digit numeric
Status	Non-core (created centrally)
Coding Source	
Coding Frame	
DD Comments	This is the unique record number used to identify each record
<hr/>	
Variable Number	1.02
Variable	Australian and New Zealand jurisdiction
Variable Name	Area
Definition	The Australian or New Zealand jurisdiction of the hospital
Justification	To enable the identification of hospitals in Australian and New Zealand jurisdictions
Format	2 digit numeric
Status	Non-core (created centrally)
Coding Source	Adapted from the National Health Data Dictionary, Version 15 (METeOR identifier 269941)
Coding Frame	1 New South Wales 2 Victoria 3 Queensland 4 South Australia 5 Western Australia 6 Tasmania 7 Northern Territory 8 Australian Capital Territory 9 Other Territories (Cocos Keeling Islands, Christmas Island and Jervis Bay Territory) 10 New Zealand
DD Comments	The order used here is the standard for the Australian Bureau of Statistics (ABS).
<hr/>	

Variable Number	1.03
Variable	First name of patient
Variable Name	Name
Definition	First name of the patient
Justification	To allow for checking of duplicate entries for the one person and to contact the patient for the 120 day follow-up
Format	Character
Status	Core
Coding Source	To allow for checking of duplicate entries for the one person as well as the ability to follow up patient including future data linkage
Coding Frame	Character string
DD Comments	The format should be the same as that indicated by the person (for example written on a form) or in the same format as that printed on an identification card, such as Medicare card, to ensure consistent collection of name data
<hr/>	
Variable Number	1.04
Variable	Surname of patient
Variable Name	Surname
Definition	Surname of the patient
Justification	To allow for checking of duplicate entries for the one person as well as the ability to follow up patient including future data linkage
Format	Character
Status	Core
Coding Source	
Coding Frame	
DD Comments	The format should be the same as that indicated by the person (for example written on a form) or in the same format as that printed on an identification card, such as Medicare card, to ensure consistent collection of name data
<hr/>	
Variable Number	1.05
Variable	Hospital MRN / URN / event number
Variable Name	MRN
Definition	Hospital Medical Record Number
Justification	Unique person-identifier for each patient in each hospital and contributes to collection of information on follow up e.g. re-operation
Format	String XXXXXX[X(14)]
Status	Core
Coding Source	
Coding Frame	
DD Comments	Key field: must be entered to create a patient record. Individual hospitals use their own alphabetic, numeric, or alphanumeric coding systems. With the eventual move to E-Health in Australia, each patient will have a unique id nation-wide. Note: Western Australia uses URN. New Zealand to record the hospital event number.
<hr/>	

Variable Number 1.06
Variable **Contact telephone number for patient**
Variable Name phone
Definition Contact telephone number of the patient
Justification To contact the patient for the 120 day follow-up
Format 10 digit numeric
Status Core
Coding Source
Coding Frame
DD Comments Only record one telephone number. This should be the best land line telephone or mobile phone number to contact the patient for the 120 day follow-up. Record the prefix plus telephone number without punctuation, for example, 08 8226 6000 or 0417 123456.

Variable Number 1.07
Variable **Date of birth**
Variable Name DOB
Definition Date of birth of the patient
Justification Basic demographic details. Required for probabilistic data linkage
Format 8 digit, date in DDMMYYYY
Status Core
Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 287007)
Coding Frame DD/MM/YYYY
DD Comments Key field Australia: must be entered to create a patient record. Only include people who are 50 years and older at the time of their hip fracture admission. Date not known is recorded as: 01011900.

Variable Number 1.08
Variable **Age derived**
Variable Name Age
Definition Age of the patient in (completed) years at admission
Justification Basic demographic details
Format 3 digit, N[NN]
Status Non-core (created centrally)
Coding Source National Health Data Dictionary, Version 15 (METeOR identifier 303794)
Coding Frame Unknown/Not stated
DD Comments If age (or date of birth) is unknown or not stated, and cannot be estimated, leave blank. Age to be calculated automatically from Date of Birth and ED/hospital arrival date (operating hospital) or ED/hospital arrival date (transfer hospital) for patients transferred to an operating hospital

Variable Number	1.09
Variable	Sex of person
Variable Name	Sex
Definition	Sex of the patient
Justification	Basic demographic details
Format	1 digit numeric
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	1 Male 2 Female 3 Intersex or indeterminate 9 Not stated / inadequately described
DD Comments	Key field: must be entered to create a patient record.

Variable Number	1.10
Variable	Australian Indigenous status
Variable Name	Indig
Definition	Was the patient of Aboriginal or Torres Strait Islander origin?
Justification	Basic demographic details
Format	1 digit numeric, N
Status	Core
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 291036)
Coding Frame	1 Aboriginal but not Torres Strait Islander origin 2 Torres Strait Islander but not Aboriginal origin 3 Both Aboriginal and Torres Strait Islander origin 4 Neither Aboriginal or Torres Strait Islander origin 9 Not stated / inadequately described
DD Comments	An Aboriginal or Torres Strait Islander is a person of Aboriginal or Torres Strait Islander descent who identifies as an Aboriginal or Torres Strait Islander and is accepted as such by the community in which he or she lives. Collected Australia only

Variable Number	1.11
Variable	NZ ethnic status
Variable Name	ethnic
Definition	Which ethnic group or groups does the patient belong to?
Justification	Basic demographic details
Format	1 digit numeric
Status	Core
Coding Source	Ministry of Health. 2017. HISO 10001:2017 Ethnicity Data Protocols. Wellington: Ministry of Health
Coding Frame	10 European 11 New Zealand European 12 Other European 21 Māori 30 Pacific peoples not further defined 31 Samoan 32 Cook Islands Māori 33 Tongan 34 Niuean 35 Tokelauan 36 Fijian 37 Other Pacific Peoples 40 Asian not further defined 41 Southeast Asian 42 Chinese 43 Indian 44 Other Asian 51 Middle Eastern 52 Latin American 53 African 61 Other Ethnicity 94 Don't Know 95 Refused to answer 97 Response unidentifiable 99 Not stated
DD Comments	<p>Patients should be asked to self-identify their ethnicity by asking them 'Which ethnic group or groups do you belong to?' For many patients it will not be possible to ask them this during their hospital admission. Therefore, the ethnicity that is recorded in the NZ hospital system should be used. The accuracy of ethnic group(s) can then be clarified at the 120 day follow up phone call. The collector must not limit the number of ethnicities given. Decisions around reporting of ethnic groups will be made in consultation with NZOA Nga Rata Koiwi representative on the NZIMC (New Zealand Implementation Committee). Collected New Zealand only.</p> <p>The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details</p>

Variable Number	1.12
Variable	Patient's postcode
Variable Name	Apcode
Definition	What was the postcode of the suburb of the usual residence of the patient?
Justification	Basic demographic details
Format	4 digit numeric, {NNNN}
Status	Core
Coding Source	Australia Post or New Zealand Post websites (www.auspost.com.au or www.nzpost.co.nz) provide up-to-date postcodes and localities
Coding Frame	1000 No fixed abode 9998 Overseas 9999 Postcode not known
DD Comments	Use a valid Australian or New Zealand postcode

Variable Number	1.13
Variable	Medicare number (Australia) / National Health Index (New Zealand)
Variable Name	Medicare
Definition	Patient's Medicare number
Justification	To allow for checking of duplicate entries for the one person and for multiple admissions
Format	Characters, N(11)
Status	Core
Coding Source	
Coding Frame	
DD Comments	<p>Enter the full Medicare number for an individual (i.e. family number plus person individual reference number).</p> <p>In Australia, if the person does not have a Medicare card or the details are unknown, enter 00000000000 (11 zeros).</p> <p>Key field New Zealand: must be entered to create a patient record. New Zealand will provide the National Health Index (NHI) which is a unique number assigned to every person who uses health and disability services in New Zealand. New Zealand will use this variable as the main mechanism to identify each patient.</p>

Variable Number	1.14
Variable	Patient type
Variable Name	ptype
Definition	Payment status
Justification	To identify the source of revenue received by a health industry relevant organisation
Format	3 digit numeric
Status	Core
Coding Source	Adapted from the National Health Data Dictionary, Version 15
Coding Frame	1 Public 2 Private 3 Overseas 9 Not known
DD Comments	<p>For New Zealand all surgery for hip fractures takes place in the public sector. There will be the occasional patient from overseas and this should be noted accordingly.</p> <p>In Australia, private sector patients include those with treatment funded by: private health insurance, workers' compensation insurance, motor vehicle third party insurance, other compensation (e.g. Public liability, common law, medical negligence), private households (i.e. self-funded and out-of-pocket expenditure), non-profit institutions serving households, corporations (other than health insurance), other private sector revenue</p> <p>In Australia, public sector patients include those with treatment funded by: Medicare, Pharmaceutical Benefits Scheme and Repatriation Pharmaceutical Benefits Scheme, Department of Veterans' Affairs, National Health and Medical Research Council, Australian Health Care Agreements, other Special Purpose payments, Other Australian Government Departments, State/Territory non-health departments, or other public sector revenue</p>

Variable Number	1.15
Variable	Usual place of residence
Variable Name	uresidence
Definition	What is the usual place of residence of the patient?
Justification	Type of accommodation before and after admission are collected to compare where the patient has come from (what was their usual accommodation) and where they are going to (what will become their usual accommodation). This is an indicator of patient outcome.
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the Australasian Rehabilitation Outcomes Centre Inpatient Dataset, Version 3.0; NSW SNAP Data Collection, Version 4.0
Coding Frame	1 Private residence (including unit in retirement village) 2 Residential aged care facility 3 Other 4 Not known
DD Comments	Record the patient's usual accommodation type at admission. Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and private hospitals or rest homes in New Zealand. If the patient lives with a relative or in a community group home or boarding house code 'private residence'. If the patient was admitted from respite care, record their usual place of residence when not in respite care.

Variable Number	1.16
Variable	Statistical linkage key 581
Variable Name	slk581
Definition	A specific code (key) that can be used to bring together two or more records belonging to the same individual. It is represented by a code consisting of characters from the person's surname, first name, date of birth and gender.
Justification	Brings together data from different sources to enable greater understanding of the utilisation of health care and/or services. Clinical quality registries should have the capacity to enhance their value through the use of linkage to other datasets (Australian Commission on Safety and Quality in Health Care Framework for Australian Clinical Quality Registries 2014)
Format	14 Characters XXXXXDDMMYYYYN
Status	Core (created centrally)
Coding Source	National Health Data Dictionary, Version 16 (METeOR identifier 349895)
Coding Frame	
DD Comments	It is represented by a code consisting of the second, third and fifth characters of a person's family name, the second and third letters of the person's given name, the day, month and year when the person was born and the sex of the person, concatenated in that order. In Australia, the linkage key is designed to make it possible to count number of clients and services they received, without counting the same client more than once. It can also be used for linking to other related data collections. This may be useful for New Zealand, although the NHI is usually the best and only identifier used for data matching in New Zealand.

Variable Number	1.17
Variable	Patient email
Variable Name	email
Definition	Email address of patient or significant other (e.g., Enduring Power of Attorney, or family member).
Justification	To contact the patient for follow up and/or to send letters and other information.
Format	String
Status	Optional, non-core
Coding Source	
Coding Frame	
DD Comments	

Variable Number	1.18
Variable	Individual Health Identifier
Variable Name	IHI
Definition	An Individual Healthcare Identifier (IHI) is a unique number used to identify an individual for health care purposes.
Justification	To enable linkage with other registries and administrative datasets such as the National Death Index and National Integrated Health Services Information (NIHSI)
Format	16 digit numeric
Status	Core
Coding Source	https://meteor.aihw.gov.au/content/432495
Coding Frame	
DD Comments	<p>Collected Australia only. All people with a Medicare card or DVA card have an IHI automatically assigned.</p> <p>New Zealand uses the National Health Index (NHI)</p> <p>New variable added 1 January 2024</p>

Section 2	Admission
------------------	------------------

Variable Number	2.01
Variable	Establishment identifier of operating hospital
Variable Name	Ahoscode
Definition	Name of the operating hospital where the patient received surgery for the hip fracture
Justification	To allow for the identification of the establishment for benchmarking and comparison purposes
Format	Character
Status	Core
Coding Source	
Coding Frame	
DD Comments	Note: For data analysis each hospital will have to be given a unique number

Variable Number	2.02
Variable	Admission via ED of operating hospital
Variable Name	EDadmit
Definition	Did the patient present directly to the ED of the operating hospital?
Justification	Ability to monitor the time spent in ED.
Format	1 digit
Status	Core
Coding Source	
Coding Frame	1 Yes 2 No - transferred from another hospital (via ED) 3 No - in-patient fall 4 No - transferred from another hospital (direct to ward) 9 Other / not known
DD Comments	If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient arrived and left the ED of the operating hospital will be recorded. If the patient was admitted directly to a ward at the operating hospital, information on the date and time that the patient arrived on the ward will be recorded. The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	2.03
Variable	.Transfer hospital
Variable Name	Athoscode
Definition	Name of the hospital where the patient first presented and was diagnosed with a hip fracture
Justification	To allow for the identification of the establishment for benchmarking and comparison purposes
Format	Character
Status	Core
Coding Source	
Coding Frame	Not transferred If transferred enter hospital name of first transfer hospital
DD Comments	If the patient has not been transferred, this field will be hidden and will not be available. Note: For data analysis, each hospital will be given a unique number. If patient is not transferred, data variables 2.04 and 2.05 regarding transfer date/time will be hidden and will not be available.

Variable Number	2.04
Variable	ED / hospital arrival date (transfer hospital)
Variable Name	tarrdate
Definition	Date on which the patient presented to the transferring hospital with a hip fracture
Justification	To enable the identification of the date of arrival in transferring hospital. Will allow for quantification of true time to surgery and overall LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	If the patient is transferred several times, this should be the hospital where the patient first presented with the hip fracture. If the presenting hospital has no ED or the patient wasn't admitted through ED, state the date presenting to the transferring hospital with a hip fracture. If the hip fracture occurred as an in-patient, record the date the fracture was diagnosed. If the date is unknown, leave blank.

Variable Number	2.05
Variable	ED arrival time (transfer hospital)
Variable Name	tarrtime
Definition	Time at which the patient arrived in the ED of the transferring hospital
Justification	To enable the identification of the time of arrival in the ED
Format	4 digit
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	hh:mm
DD Comments	Time is recorded using the 24 hour clock.

If the patient is transferred several times, this should be the hospital where the patient first presented with a hip fracture.

If the presenting hospital has no ED or the patient wasn't admitted through ED, state the time presenting to the transferring hospital with a hip fracture.

If the hip fracture occurred as an in-patient, record the time the fracture was diagnosed. Note: 00:00 indicates that the time was not known.

To be used in the calculation of total LOS in the hospital system for the care episode.

Variable Number	2.06
Variable	ED / other ward arrival date (operating hospital)
Variable Name	arrdate
Definition	Date on which the patient arrived in the ED / other ward of the operating hospital
Justification	To enable calculation of age at presentation, time spent in ED, time to surgery and LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	<p>If the patient was not admitted through the ED but was transferred from another hospital and admitted directly to a ward of the operating hospital, state the date admitted to the ward of the operating hospital.</p> <p>If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded. The Australian National Emergency Access Target (NEAT) aims that by 2015, 90% of patients will leave the ED within 4 hours – either by discharge, being admitted to hospital or through transfer to another hospital for treatment (http://www.ecinsw.com.au/node/128). For New Zealand patients are expected to be discharged or admitted to hospital within 6 hours.</p>

Variable Number	2.07
Variable	ED / other ward arrival time (operating hospital)
Variable Name	arrtime
Definition	Time at which the patient arrived at the ED / other ward of the operating hospital
Justification	To enable calculation of time spent in ED, time to surgery and LOS
Format	5 digit
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	hh:mm
DD Comments	Time is recorded using the 24 hour clock.

If the patient was not admitted through the ED but was transferred from another hospital and admitted directly to a ward of the operating hospital, state the time admitted to the ward of the operating hospital.

Note: 00:00 indicates that the time was not known.

If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded.

Variable Number	2.08
Variable	ED departure date (operating hospital)
Variable Name	depdate
Definition	Date on which the patient departed from the ED of the operating hospital
Justification	To enable calculation of time spent in ED, time to surgery and LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	Note: If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded

Variable Number	2.09
Variable	ED departure time (operating hospital)
Variable Name	deptime
Definition	Time at which the patient departed from the ED of the operating hospital
Justification	To enable calculation of time spent in ED, time to surgery and LOS
Format	4 digit
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	hh:mm
DD Comments	Time is recorded using the 24 hour clock.

Note: 00:00 indicates that the time was not known.

If the patient was admitted via the ED of the operating hospital, information on the date and time that the patient left the ED of the operating hospital will be recorded.

Variable Number	2.10
Variable	In-patient fracture date
Variable Name	admdateop
Definition	Date on which the admitted patient commences the episode of care at the operating hospital with radiological-confirmed diagnosis of hip fracture
Justification	To enable the identification of the date of hip fracture occurring as an in-patient and calculation of time to surgery and LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	If the date is unknown, leave blank. Fractures sustained while on leave from an existing hospital admission are not classified as inpatient fractures. They are recorded as a new event and date and time of presentation are recorded at 2.06 and 2.07.

Variable Number	2.11
Variable	In-patient fracture time
Variable Name	admtimeop
Definition	24-hour time at which the admitted patient commences the episode of care at the operating hospital with radiological-confirmed diagnosis of hip fracture
Justification	To enable the identification of the time of hip fracture occurring as an in-patient and calculation of time to surgery and LOS
Format	4 digit
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	hh:mm
DD Comments	Time is recorded using the 24 hour clock. Note: 00:00 indicates that the time was not known.

Variable Number	2.13
Variable	Pain management
Variable Name	painmanage
Definition	Did the patient receive analgesia within 30 minutes of presentation to the emergency department?
Justification	Acute pain associated with the hip fracture can have adverse effects on outcome. Hip Fracture Clinical Care Standard Indicator 2b.
Format	1 digit
Status	Core
Coding Source	
Coding Frame	1 Analgesia given within 30 minutes of ED presentation 2 Analgesia given more than 30 minutes after ED presentation 3 Analgesia not required – already provided by paramedics 4 Analgesia not required – no pain documented on assessment 9 Not known
DD Comments	Time to analgesia in the ED to be identified from clinical notes. Time is calculated from date and time of presentation to the emergency department of the first hospital. Coding frame option 3 includes provision of analgesia by paramedics at the scene, in-transit and after arrival to hospital. This is used in cases where the patient does not need or is not due for additional analgesia within 30 minutes of arrival to ED. New variable added 1 January 2017. The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	2.15
Variable	.Transferred patients only: Nerve block before transfer
Variable Name	tfanalges
Definition	Did the patient with a hip fracture who was transferred from another hospital for treatment receive a nerve block prior to transfer?
Justification	Hip Fracture Clinical Care Standard Indicator 2b
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	1 No 2 Yes 9 Not known
DD Comments	If the patient was not transferred, record information on nerve blocks under variable 4.06 Analgesia-nerve block

Variable Number	3.01
Variable	Pre-admission walking ability
Variable Name	walk
Definition	What was the patient's walking ability pre-admission?
Justification	To enable the identification of the mobility status pre-admission
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Usually walks without walking aids 2 Usually walks with either a stick or crutch 3 Usually walks with two aids or frame (with or without assistance of a person) 4 Usually uses a wheelchair / bed bound 9 Not known
DD Comments	<p>If a person has different levels of mobility on different surfaces then record the level of most assistance. For example, inside their residence a person usually walks without a walking aid but when outside the residence the person usually walks with a frame, then the level of mobility recorded is option 3.</p> <p>If a person does not use a walking aid, but requires physical assistance e.g. from a carer or staff in Residential care, pre-admission walking ability should be coded as following: 2 if assistance of one person is required and; 3 if assistance of two people is required</p>

Variable Number	3.02
Variable	Pre-operative cognitive assessment
Variable Name	cogassess
Definition	Following admission to hospital, cognitive status is assessed prior to surgery using a validated tool and recorded in the medical record
Justification	Hip fracture patients are at high risk of having an existing cognitive impairment or developing delirium. Cognitive impairment and delirium in these patients is associated with increased morbidity and mortality, and a decrease in rehabilitation potential and return to pre-fracture functioning. Care at Presentation Hip Fracture Clinical Care Standard Indicator 1a.
Format	1 digit
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Not assessed 2 Assessed and normal 3 Assessed and abnormal or impaired 9 Not known
DD Comments	<p>Cognitive assessment requires the use of a validated tool. Some validated tools for assessing cognitive function include:</p> <ul style="list-style-type: none"> • Abbreviated Mental Test Score (AMTS) (Hodkinson 1972) • Standardised Mini-Mental State Examination (SMMSE) (Molloy & Standish 1997) • Modified Mini Mental State Exam (3MS) (Teng & Chui 1987) • General Practitioner's Assessment of Cognition (GPCOG) (Brodady et al. 2002) • The 4AT (Bellelli et al. 2014) • Other tools, such as the Rowland Universal Dementia Assessment Scale (RUDAS) (Storey et al. 2004) and the Kimberly Indigenous Cognitive Assessment (KICA) (LoGiudice et al. 2006), may be more appropriate for some people from culturally and linguistically diverse groups <p>New variable added 1 January 2017. The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details</p>

Variable Number	3.05
Variable	Pre-admission cognitive status
Variable Name	cogstat
Definition	What was the cognitive status of the patient prior to admission?
Justification	To enable the identification of the cognitive status of the patient prior to admission.
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	1 Normal cognition 2 Impaired cognition or known dementia 9 Not known
DD Comments	Normal cognition refers to ‘no history of cognitive impairment or dementia’. Impaired cognition or known dementia refers to a ‘loss of cognitive ability and/or a decline in memory or other thinking skills severe enough to reduce a person’s ability to perform everyday activities’ (Alzheimer’s Association).
	The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details


Variable Number	3.06
Variable	Bone protection medication at admission
Variable Name	bonemed
Definition	Was the patient taking bone protection medication prior to sustaining the hip fracture?
Justification	Ability to monitor use of bone protection medication prior to hip fracture
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No bone protection medication 1 Calcium and/or vitamin D only 2 Yes - bisphosphonates, denosumab, romosozumab, teriparatide, raloxifene or HRT (with or without calcium and/or vitamin D) 9 Not known
DD Comments	Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or one alpha).
	Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate, Zoledronate, Pamidronate.
	A patient is to be considered as taking osteoporosis specific treatment if: <ul style="list-style-type: none"> • Oral bisphosphonates, prescribed in the last 12 weeks. • Zoledronate, administered in the last 24 months. • Denosumab, administered the last 6 months. • Teriparatide, administered in the last 7 days. • Romosozumab, administered in the last month. These medications may be prescribed with or without calcium and / or vitamin D.
	For changes to this variable over time, please see current Data Variable Concordance table at https://anzhfr.org/data-access/

Variable Number	3.07
Variable	Pre-operative medical assessment
Variable Name	passess
Definition	Who conducted the pre-operative medical assessment apart from anaesthetic and orthopaedic review?
Justification	To determine level of pre-operative medical assessment.
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No assessment conducted 1 Geriatrician / Geriatric Team 2 Physician / Physician Team 3 GP 4 Specialist nurse 9 Not known
DD Comments	The pre-operative assessment is conducted in addition to an anaesthetic review and orthopaedic assessment. If the pre-operative assessment is conducted by a number of assessment team members, select the highest numerical option in the coding frame drop down list eg. the highest numerical option to select is '1' geriatrician.

Variable Number	3.08
Variable	Side of hip fracture
Variable Name	side
Definition	What was the side of the patient's hip fracture?
Justification	To enable the identification of the side of the hip fracture
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	1 Left 2 Right
DD Comments	Key field: must be entered to create a patient record. If the patient has bilateral hip fractures, a separate record should be created for each fracture.

Variable Number	3.09
Variable	Atypical fracture
Variable Name	afraction
Definition	Was the type of the patient's hip fracture either pathological or atypical?
Justification	To enable the identification of fractures which are not consistent with the nature of the injury
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 Not a pathological or atypical fracture 1 Pathological fracture 2 Atypical fracture
DD Comments	A pathological fracture is considered to be a fracture that has occurred when a bone breaks in an area that has been weakened by another disease process (except osteoporosis), such as a tumour, infection or an inherited bone disorder. An atypical fracture is one where the radiologically observed fracture pattern is not consistent with the mechanism of injury described and is not thought to be attributable to a discrete underlying disease process

Variable Number	3.10
Variable	.Type of fracture
Variable Name	ftype
Definition	What was the type of the patient's hip fracture?
Justification	To enable the identification of the type of hip fracture
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Intracapsular undisplaced/impacted displaced 2 Intracapsular displaced 3 Per/intertrochanteric 4 Subtrochanteric
DD Comments	Basal/basicervical fractures are to be classified as per/intertrochanteric



Variable Number	3.11
Variable	.Surgical repair
Variable Name	surg
Definition	Did the patient undergo surgical repair of the hip fracture?
Justification	To enable quantification of percentage patients undergoing surgery
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	2 Yes 3 No – surgical fixation not clinically indicated 4 No – patient for palliation 5 No – other reason
DD Comments	The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	3.12
Variable	ASA grade
Variable Name	asa
Definition	What is the ASA grade for the patient?
Justification	A marker of disease severity and operative risk and used for case-mix adjustment
Format	1 digit numeric
Status	Core
Coding Source	American Society of Anaesthesiologists
Coding Frame	1 Healthy individual with no systemic disease 2 Mild systemic disease not limiting activity 3 Severe systemic disease that limits activity but is not incapacitating 4 Incapacitating systemic disease which is constantly life threatening 5 Moribund not expected to survive 24 hours with or without surgery 9 Not known
DD Comments	ASA grade is used in case-mix adjustment for outcome at 30 and 120 days post-surgery

Variable Number	3.13
Variable	Clinical Frailty Scale
Variable Name	frailty
Definition	What was the patient's pre-injury frailty status?
Justification	To enable the identification of the patient's frailty status prior to their hip fracture as a person's level of frailty impacts outcomes. Hip Fracture Clinical Care Standard Indicator 3a.
Format	2 digit numeric
Status	Core
Coding Source	Rockwood Clinical Frailty Scale
Coding Frame	1 Very Fit 2 Well 3 Well, with treated comorbid disease 4 Vulnerable 5 Mildly frail 6 Moderately frail 7 Severely frail 8 Very severely frail 9 Terminally ill 10 Frailty assessment using other validated tool 99 Not known
DD Comments	NOTE: the Clinical Frailty Scale applies to the person's usual status prior to the hip fracture. Where the person has dementia or delirium the information will need to be provided by an informant who knows the person well.

Coding Frame Definitions

- 1 **Very fit** - robust, active, energetic and well-motivated. Exercise regularly and are among the fittest for their age.
- 2 **Well** - without active disease symptoms but are less fit than category 1. Exercise occasionally.
- 3 **Well with treated comorbid disease** - disease symptoms are well controlled compared to category four. Not regularly active beyond routine walking.
- 4 **Vulnerable** - not dependent on others for daily help, but symptoms limit activities. Common complaint is being 'slowed up' or being tired during the day.
- 5 **Mildly frail** - more evident slowing, and need help in instrumental activities of daily living (e.g. heavy housework, medications, transportation, shopping, using the phone, managing finances, meal preparation).
- 6 **Moderately frail** - need help with both instrumental and non-instrumental activities of daily living. Includes mobility in bed, transferring on/off chairs, toilets and into/out of bed, walking, dressing, eating, toilet use, personal hygiene, bathing.
- 7 **Severely frail** - completely dependent on others for all activities of daily living for whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~ 6 months).
- 8 **Very severely frail** - completely dependent on others for all activities of daily living, approaching the end of life. Typically, they could not recover even from a minor illness.
- 9 **Terminally ill** - approaching the end of life. Applies to people with a life expectancy <6 months who are not otherwise evidently frail.

New variable added 1 January 2021.

The coding frame for this variable has changed. Please see current Data Variable Concordance table at <https://anzhfr.org/data-access/> for details

Variable Number	3.14
Variable	Delirium assessment prior to surgery
Variable Name	addelassess
Definition	Following admission to hospital, was delirium assessed prior to surgery using a validated tool and recorded in the medical record?
Justification	Identifying patients with delirium is the first step in taking action to providing high quality care. Early diagnosis and prompt treatment offers patients with delirium the best chance of recovery.
Format	1 digit
Status	Non-Core
Coding Source	
Coding Frame	1 Not assessed 2 Assessed and not identified 3 Assessed and identified 9 Not known
DD Comments	<p>Assessment of delirium requires the use of a validated tool. There are a range of validated diagnostic tools for delirium and they include:</p> <ul style="list-style-type: none"> • Confusion Assessment Method (CAM) (Inouye et al. 2014; Shi et al. 2013) • Confusion Assessment Method (CAM-ICU) (Ely et al. 2001) • 3D-CAM (Marcantonio et al. 2014). • The 4AT (Bellelli et al. 2014) <p>If a person declines assessment record as not assessed.</p> <p>Delirium is defined as an acute change in mental status that is common among older patients in hospital (Clinical Epidemiology and Health Service Evaluation Unit 2006). It is characterised by a disturbance of consciousness, attention, cognition and perception that develops over a short period of time (usually hours to a few days) (National Institute for Health and Clinical Excellence 2010; Inouye 2014). Patients with delirium may be agitated and restless (hyperactive delirium), quiet and withdrawn (hypoactive delirium), or move between these two subtypes (mixed delirium) (Clinical Epidemiology and Health Service Evaluation Unit 2006; National Institute for Health and Clinical Excellence 2010).</p> <p>New variable added 1 January 2024. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/</p>

Section 4	Treatment
------------------	------------------

Variable Number 4.01
Variable .Date of surgery for hip fracture
Variable Name sdate
Definition Date on which the surgery for the hip fracture takes place
Justification To enable the identification of the date of primary surgery. Hip Fracture Clinical Care Standard Indicator 4a.
Format 8 digit date, date in DDMMYYYY
Status Core
Coding Source National Health Data Dictionary, Version 15
Coding Frame DD/MM/YYYY
DD Comments If the date is unknown, leave blank.

Variable Number 4.02
Variable .Time of surgery for hip fracture
Variable Name stime
Definition 24-hour time at which the surgery for the hip fracture commences. This time is taken from the start of the anaesthetic process.
Justification To enable the identification of the start time of the primary surgery. Hip Fracture Clinical Care Standard Indicator 4a.
Format 4 digit
Status Core
Coding Source National Health Data Dictionary, Version 15
Coding Frame hh:mm
DD Comments Time is recorded using the 24 hour clock

The time of surgery for the hip fracture is taken from the start of the anaesthetic process.

Note: 00:00 indicates that the time was not known.

Variable Number	4.03
Variable	Surgery delay
Variable Name	delay
Definition	What was the primary reason for the delay if the delay was greater than 36 hours from the time of arrival in the emergency department of the first hospital, or diagnosis of a fracture if the fracture occurred as an in-patient?
Justification	Ability to monitor time to surgery as a standard of care. Hip Fracture Clinical Care Standard Indicator 4a.
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	2 Delay due to patient deemed medically unfit 3 Delay due to issues with anticoagulation 4 Delay due to theatre availability 5 Delay due to surgeon availability 6 Delay due to delayed diagnosis of hip fracture 7 Other type of delay (state reason) 9 Not known
DD Comments	<p>Delay is calculated from the time of presentation in the emergency department of the first hospital.</p> <p>A person is considered medically unfit if he/she have acute health-related issues which need to be stabilised/optimised or reversed prior to proceeding with anaesthesia and a surgical procedure.</p> <p>If there is more than one delay to surgery, choose the reason for the first delay.</p> <p>The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details</p>

Variable Number	4.04
Variable	Surgery delay other text
Variable Name	delay_txt
Definition	What was the reason for the other delay, if the delay was greater than 36 hours from the time of arrival in the emergency department?
Justification	Ability to monitor time to surgery as a standard of care
Format	Character
Status	Core
Coding Source	
Coding Frame	
DD Comments	

Variable Number	4.05
Variable	.Type of anaesthesia
Variable Name	anaesth
Definition	What type of anaesthesia for the hip fracture surgery?
Justification	Ability to monitor variation, post-operative complications and patient choice
Format	2 digit numeric
Status	Core
Coding Source	
Coding Frame	1 General anaesthesia 2 Spinal anaesthesia 3 General and spinal anaesthesia 97 Other 99 Not known
DD Comments	CSE=Combined Spinal/Epidural. Recorded in anaesthetic chart Information on administering a peripheral nerve block is recorded in 'Analgesia – nerve block'. The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	4.06
Variable	.Analgesia - nerve block
Variable Name	analges
Definition	Did the patient have a nerve block?
Justification	Hip Fracture Clinical Care Standard Indicators 2b and 2c
Format	2 digit numeric
Status	Core
Coding Source	
Coding Frame	1 Nerve block administered before arriving in OT 2 Nerve block administered in OT 3 Both 4 Neither 99 Not known
DD Comments	

Variable Number	4.07
Variable	Consultant surgeon present
Variable Name	consult
Definition	Was the consultant surgeon operating or assisting with the operation?
Justification	Ability to monitor the impact of consultant surgeon presence on the quality and safety of patient outcome
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	0 No 1 Yes 9 Not known
DD Comments	To record yes, consultant must be scrubbed and operating. This variable can be found by checking if the consultant surgeon is recorded on the operation sheet
	The DD comments for this variable have changed over time. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	4.08
Variable	Type of operation performed
Variable Name	optype
Definition	What type of operation was performed for the hip fracture?
Justification	To enable the identification of the patient's type of hip fracture operation
Format	2 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Cannulated screws (e.g. multiple screws) 2 Sliding hip screw 3 Intramedullary nail short 4 Intramedullary nail long 5 Hemiarthroplasty stem cemented 6 Hemiarthroplasty stem uncemented 7 Total hip replacement stem cemented 8 Total hip replacement stem uncemented 9 Femoral neck system (FNS) 97 Other 99 Not known
DD Comments	iFor cemented versus uncemented procedures, this only includes whether the stem was cemented or not. This does not include whether or not the cup was cemented. Sliding hip screws include dynamic hip screws (DHS). The Femoral Neck System may also be referred to as the DePuy Synthes Femoral Neck System.

Variable Number	4.10
Variable	Full weight bear
Variable Name	wbear
Definition	What is the patient's immediate post-operative weight bearing status?
Justification	Ability to monitor variation in practice. Hip Fracture Clinical Care Standard Indicator 5b.
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	0 Unrestricted weight bearing 1 Restricted / non weight bearing 9 Not known
DD Comments	Unrestricted weight bearing refers to a patient who is able to mobilise with full use of the affected limb to weight bear as pain allows. Restricted weight bearing refers to a patient where there is a specific instruction that prevents the patient being allowed to fully utilise the leg irrespective of degree of pain. Restricted weight bearing includes terms such as partial weight bear, touch-weight bear and non-weight bear
Variable Number	4.12
Variable	New pressure injuries of the skin
Variable Name	Pulcers
Definition	Did the patient acquire a new pressure injury (Stage II or above) during their stay in hospital for the treatment of their hip fracture?
Justification	Hip Fracture Clinical Care Standard Indicator 5b Pressure injuries of the skin are potentially preventable. They can affect a person's level of pain, quality of life, cost of care, and mortality.
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No 1 Yes 9 Not known
DD Comments	A pressure injury is an area of localised damage to the skin and underlying tissue caused by pressure, shear or friction forces, or a combination of these. Grading for pressure ulcers consists of 4 levels: Stage I pressure injury: non-blanchable erythema (intact skin with non-blanchable redness of a localised area usually over a bony prominence). Stage II pressure injury: partial thickness skin loss (Partial thickness loss of dermis presenting as a shallow open ulcer with a red pink wound bed, with slough). Stage III pressure injury: full thickness skin loss (Subcutaneous fat may be visible but bone, tendon, or muscle, are not fully exposed). Stage IV pressure injury: full thickness tissue loss (Full thickness tissue loss with exposed bone, tendon or muscle). The pressure injury classification is from the National Pressure Ulcer Advisory Panel (NPUAP) and European Pressure Ulcer Advisory Panel (EPUAP), Prevention and Treatment of Pressure Ulcers: Clinical Practice Guideline. 2009, Washington DC: NPUAP

Variable Number	4.13
Variable	Assessed by geriatric medicine
Variable Name	gerimed
Definition	Was the patient assessed by geriatric medicine during the acute phase of the episode of care?
Justification	Ability to monitor quality of care. Hip Fracture Clinical Care Standard Indicator 3a.
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	0 No 1 Yes 8 No geriatric medicine service available 9 Not known
DD Comments	An assessment by geriatric medicine refers to an assessment by a geriatrician or a medical practitioner (Registrar) working under the supervision of a geriatrician. The acute phase (IHPA Admitted Hospital Care Types: Guide For Use 2015) is care in which the primary clinical purpose or treatment goal is to: <ul style="list-style-type: none"> • cure illness or provide definitive treatment of injury • perform surgery • relieve symptoms of illness or injury (excluding palliative care) • reduce severity of an illness or injury • protect against exacerbation and/or complication of an illness and/or injury which could threaten life or normal function • perform diagnostic or therapeutic procedures

Variable Number	4.14
Variable	Geriatric medicine assessment date
Variable Name	gdate
Definition	Date on which an admitted patient was first assessed by geriatric medicine during the acute phase of their episode of care
Justification	To enable the identification of the date of geriatric assessment. Hip Fracture Clinical Care Standard Indicator 3a.
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	A geriatric assessment is considered to include an assessment by a geriatrician or a medical practitioner (Registrar) working under the supervision of a geriatrician. Leave blank if not known



Variable Number	4.16
Variable	Bone protection medication at discharge from hospital
Variable Name	dbonemed1
Definition	What bone protection medication was the patient taking at discharge from hospital?
Justification	Ability to monitor use of bone protection medication. Hip Fracture Clinical Care Standard Indicator 6a.
Format	1 digit numeric
Status	Code
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No bone protection medication 1 Calcium and/or vitamin D only 2 Yes - Bisphosphonates, denosumab, romosozumab, teriparatide, raloxifene or HRT (with or without calcium and/or vitamin D) 3 No but received prescription at separation from hospital 9 Not known
DD Comments	<p>Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or one alpha).</p> <p>Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate, Zoledronate, Pamidronate.</p> <p>A patient is to be considered as taking osteoporosis specific treatment if:</p> <ul style="list-style-type: none"> • Oral bisphosphonates, prescribed in the last 12 weeks. • Zoledronate, administered in the last 24 months. • Denosumab, administered the last 6 months. • Teriparatide, administered in the last 7 days. • Romosozumab, administered in the last month. <p>These medications may be prescribed with or without calcium and / or vitamin D.</p> <p>For changes to this variable over time, please see current Data Variable Concordance table at https://anzhfr.org/data-access/</p>

Variable Number	4.17
Variable	Post-operative delirium assessment
Variable Name	delassess
Definition	Did the patient have a documented assessment of delirium in the week following surgery for their hip fracture?
Justification	Hip Fracture Clinical Care Standard Indicator 3b Identifying patients with delirium is the first step in taking action to providing high quality care. Early diagnosis and prompt treatment offers patients with delirium the best chance of recovery.
Format	1 digit
Status	Non-Core
Coding Source	
Coding Frame	1 Not assessed 2 Assessed and not identified 3 Assessed and identified 9 Not known
DD Comments	<p>Assessment of delirium requires the use of a validated tool. There are a range of validated diagnostic tools for delirium and they include:</p> <ul style="list-style-type: none"> • Confusion Assessment Method (CAM) (Inouye et al. 2014; Shi et al. 2013) • Confusion Assessment Method (CAM-ICU) (Ely et al. 2001) • 3D-CAM (Marcantonio et al. 2014). • The 4AT (Bellelli et al. 2014) <p>If a person declines assessment record as not assessed.</p> <p>Delirium is defined as an acute change in mental status that is common among older patients in hospital (Clinical Epidemiology and Health Service Evaluation Unit 2006). It is characterised by a disturbance of consciousness, attention, cognition and perception that develops over a short period of time (usually hours to a few days) (National Institute for Health and Clinical Excellence 2010; Inouye 2014). Patients with delirium may be agitated and restless (hyperactive delirium), quiet and withdrawn (hypoactive delirium), or move between these two subtypes (mixed delirium) (Clinical Epidemiology and Health Service Evaluation Unit 2006; National Institute for Health and Clinical Excellence 2010).</p> <p>New variable added 1 January 2018. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/</p>

Variable Number	4.18
Variable	Clinical malnutrition assessment
Variable Name	malnutrition
Definition	Did the patient undergo clinical assessment of their protein/energy nutrition status during the acute phase of the episode of care?
Justification	Hip fracture patients are at high risk of malnutrition. Malnutrition in these patients is associated with increased morbidity and mortality, and a decrease in return to pre-fracture functioning.
Format	1 digit
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 Not done 1 Malnourished 2 Not malnourished 9 Not known
DD Comments	Clinical assessment of a person's nutritional status is encouraged during the acute phase. Sites should use tools that are validated for such purposes, and are advised to discuss with their Dietitians how best to record the results using this variable's options. If the nutritional assessment is performed more than once, please record the first assessment after admission that uses a validated tool. New variable added 1 January 2019. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/

Variable Number	4.19
Variable	First day walking
Variable Name	mobil2
Definition	Did the patient get out of bed and walk on day one post hip fracture surgery?
Justification	Hip Fracture Clinical Care Standard Indicator 5a. Early mobilisation after hip fracture surgery is associated with survival and recovery (Goubar et al. 2021).
Format	1 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Yes 2 No – Stood without stepping / walking 3 No – Sat on the edge of the bed 4 No – Sat out of bed via hoist 5 No – Did not attempt to get out of bed on day one 9 Not known
DD Comments	Day 1 post-surgery means the next calendar day following the day of the patient's primary surgery for hip fracture. This data item is recording whether the patient actually stood and stepped or walked by day 1 post-surgery. Yes means the patient managed to stand and step transfer out of bed onto a chair/commode or walk. This does not include only sitting on the edge of the bed or standing up from the bed without stepping/walking. Coding frame option 2 should be selected where the patient stood with assistance but was unable to step / walk or where a standing transfer aid was used. This includes devices such as Sara Steady, Romedic Return, Standing Hoist. Coding frame option 3 should be used where the patient sat on the edge of the bed, but was not able to stand or be transferred to sitting out of bed.

Coding frame option 4 should be selected where a hoist is used to transfer the patient out of bed.

Coding frame option 5 should be used where the patient did not attempt to get out of bed.

Goubar A, Martin FC, Potter C, Jones GD, Sackley C, Ayis S, Sheehan KJ. The 30-day survival and recovery after hip fracture by timing of mobilization and dementia : a UK database study. Bone Joint J. 2021 Jul;103-B(7):1317-1324. doi: 10.1302/0301-620X.103B7.BJJ-2020-2349.R1. PMID: 34192935; PMCID: PMC7611209.

New variable added in 2020. Coding change in 2025. Please see current Data Variable Concordance table at <https://anzhfr.org/data-access/>

Variable Number	4.20
Variable	Oral nutritional supplements
Variable Name	ONS
Definition	Did the patient receive protein and energy oral nutritional supplements (ONS) during their admission?
Justification	Hip Fracture Clinical Care Standard Indicator 3c.
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	0 No 1 Yes 9 Not known
DD Comments	<p>ONS are defined as protein and energy nutrient-dense products purposed to increase dietary intake when diet alone is likely to be inadequate to meet nutritional requirements. These may include energy and protein enriched drinks (e.g. milk, soy, protein-fortified juice flavours), powders, soups, and/or desserts.</p> <p>Documented patient-level evidence must be observed as evidence of receiving ONS. Examples include documentation in the patient’s medical record (e.g. ‘patient provided with high protein drink’) or patient-specific food services system/menu orders, documentation in the medication chart or food and fluid charts.</p> <p>New variable added 1 January 2024.</p>

Variable Number	5.01
Variable	Discharge date from acute ward
Variable Name	wdisch
Definition	Date on which the patient was discharged from an acute ward during their episode of care
Justification	To enable the identification of the date of discharge from an acute ward so as to calculate LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	The discharge date refers to the patient physically leaving the acute ward. Record the date the patient was physically discharged from the acute orthopaedic stay. If the date is unknown, leave blank.
Variable Number	5.02
Variable	Discharge destination from acute ward
Variable Name	wdest
Definition	What is the discharge destination of the patient from the acute ward?
Justification	To assess patient outcome
Format	2 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Private residence (including unit in retirement village) 2 Residential aged care facility 3 Rehabilitation unit public 4 Rehabilitation unit private 5 Other hospital / ward / specialty 6 Deceased 7 Short term care in residential care facility (New Zealand only) 97 Other 99 Not known
DD Comments	Record the patient's discharge destination at discharge from the acute orthopaedic stay. If the patient is discharged to live with a relative or in a community group home or boarding house code 'private residence'. Rehabilitation unit public includes Geriatric Evaluation and Management (GEM) units. Private rehabilitation units will not be applicable in New Zealand. Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and private hospitals or rest homes in New Zealand. Short-term care in residential care facility may be relevant if the patient is non-weight bearing, and is used in New Zealand and, to a lesser degree, in Australia. For example, in New Zealand, some District Health Boards have programmes that fund patients who are non-weight bearing to be cared for in a residential care facility. When they are approved by the Orthopaedic Department to fully weight bear they are either admitted to Older Peoples Health for rehabilitation or discharged home.

Variable Number	5.03
Variable	.Discharge from hospital date
Variable Name	hdisch
Definition	Final date of discharge from an inpatient facility (public or private) after hip fracture
Justification	To enable the identification of the final date of discharge from hospital and calculation of total inpatient LOS
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	This refers to the date the patient physically leaves hospital after their hip fracture. When a person has been discharged home or to a residential aged care facility as a virtual inpatient, enter the date the patient leaves hospital. Discharge from hospital date will be the same as discharge from acute ward if patient was discharged home, to residential care or died on the acute ward. If the date is unknown, leave blank.

Variable Number	5.04
Variable	.Length of stay (acute ward)
Variable Name	olos
Definition	The length of stay of a patient in the acute ward, excluding leave days or days before fracture if occurred in hospital, measured in days
Justification	To enable the identification of the length of stay in the acute ward.
Format	3 digit numeric
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	NNN
DD Comments	Formula: Length of Stay (LOS) = Separation date - Admission date - Total leave days. The calculation is inclusive of admission and separation dates. LOS will be calculated automatically from the acute ward separation and admission dates. If the hip fracture occurred as an in-patient then the length of stay should be from time hip fracture was diagnosed.

Variable Number	5.05
Variable	Length of stay (hospital system)
Variable Name	TLOS
Definition	The length of stay of a patient from admission/diagnosis of a hip fracture to final date of discharge from an inpatient facility (public or private), excluding leave days, measured in days
Justification	To enable the identification of the total length of stay in the hospital system
Format	4 digit, unit of measure (day)
Status	Non-core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	NNNN
DD Comments	<p>Formula: Length of stay (LOS) = Separation date - Admission date - Total leave days. The calculation is inclusive of admission and separation dates.</p> <p>LOS will be calculated automatically from the ED arrival date of the transferring hospital (or the ED arrival date of the operating hospital, if no transfer occurred) and the discharge from hospital date. If the hip fracture occurred as an in-patient then the length of stay should be from time hip fracture was diagnosed. If the final date of discharge from the hospital system is known, this date should be used.</p> <p>It should be noted that the total length of stay in the hospital system will be difficult to calculate in some jurisdictions, due to differences in treatment settings for rehabilitation-based care.</p>

Variable Number	5.06
Variable	Discharge place of residence
Variable Name	dresidence
Definition	What is the usual place of residence of the person following discharge from the whole hospital system?
Justification	Type of accommodation before and after admission are collected to compare where the patient has come from (what was their usual accommodation) and where they are going to (what will become their usual accommodation). Comparison of accommodation pre and post admission is an indicator of patient outcome.
Format	1 digit numeric
Status	Core
Coding Source	Adapted from Australasian Rehabilitation Outcomes Centre Inpatient Dataset, Version 3.0; NSW SNAP Data Collection, Version 4.0
Coding Frame	1 Private residence (including unit in retirement village) 2 Residential aged care facility 3 Deceased 7 Other 9 Not known
DD Comments	Record the patient's accommodation type at discharge from the whole hospital system. If the patient lives with a relative or in a community group home or boarding house code 'private residence'. Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and private hospitals or rest homes in New Zealand.

Section 7	120 day follow-up*
------------------	---------------------------

*120-day follow up is undertaken by the operating hospital

Variable Number	7.01
Variable	.120 day follow-up date
Variable Name	fdate2
Definition	Date on which the 120 day follow-up was completed post the initial hip fracture surgery
Justification	To monitor patient outcomes post-surgery
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	Date not known is left blank

Variable Number	7.02
Variable	.Survival at 120 days post-surgery
Variable Name	fsurvive2
Definition	Is the patient alive at 120 days post-surgery
Justification	To monitor patient outcomes post-surgery
Format	1 digit numeric
Status	Core
Coding Source	
Coding Frame	0 No 1 Yes 9 Not known
DD Comments	

Variable Number	7.03
Variable	.Date hospital system discharge at 120 day follow-up
Variable Name	date120
Definition	What date was the patient discharged from the hospital system?
Justification	To enable the identification of the total length of stay in the hospital system
Format	8 digit date, date in DDMMYYYY
Status	Core
Coding Source	National Health Data Dictionary, Version 15
Coding Frame	DD/MM/YYYY
DD Comments	

Variable Number	7.04
Variable	Place of residence at 120 day follow-up
Variable Name	fresidence2
Definition	What is the place of residence of the person at 120 days post-surgery?
Justification	To monitor patient outcomes post-surgery. Hip Fracture Clinical Care Standard Indicator 7b.
Format	2 digit numeric
Status	Core
Coding Source	Adapted from the Australasian Rehabilitation Outcomes Centre Inpatient Dataset, Version 3.0; NSW SNAP Data Collection, Version 4.0
Coding Frame	<ul style="list-style-type: none"> 1 Private residence (including unit in retirement village) 2 Residential aged care facility 3 Rehabilitation unit public 4 Rehabilitation unit private 5 Other hospital / ward / specialty 6 Deceased 7 Short term care in residential care facility (New Zealand only) 97 Other 99 Not known
DD Comments	<p>Record the patient's discharge destination at 120 days post-surgery. If the patient is discharged to live with a relative or in a community group home or boarding house code 'private residence'. Private rehabilitation units will not be applicable in New Zealand.</p> <p>Residential aged care facility refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and private hospitals or rest homes in New Zealand.</p> <p>Short-term care in residential care facility may be relevant if the patient is non-weight bearing and is used in New Zealand and to a lesser degree in Australia. For example, in New Zealand, some District Health Boards have programmes that fund patients who are non-weight bearing to be cared for in a residential care facility. When they are approved by the Orthopaedic Department to fully weight bear they are either admitted to Older Peoples Health for rehabilitation or discharged home.</p>

Variable Number	7.06
Variable	Post-admission walking ability at 120 day follow-up
Variable Name	fwalk2
Definition	What was the patient's walking ability at 120 days post-surgery?
Justification	To monitor patient mobility status post-discharge. Hip Fracture Clinical Care Standard Indicator 5d.
Format	2 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	1 Usually walks without walking aids 2 Usually walks with either a stick or crutch 3 Usually walks with two aids or frame 4 Usually uses a wheelchair / bed bound 8 Not relevant 9 Not known
DD Comments	Usually walks with two aids or frame includes with or without assistance of a person If a person has different levels of mobility on different surfaces then record the level of most assistance. For example, inside their residence a person usually walks with no walking aid but when outside the residence the person usually walks with a frame, then the level of mobility recorded is option 3. The coding frame for this variable has changed. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/ for details

Variable Number	7.07
Variable	Bone protection medication at 120 day follow-up
Variable Name	fbonemed2
Definition	What bone protection medication was the patient using at 120 days post-surgery?
Justification	Ability to monitor use of bone protection medication
Format	1 digit numeric
Status	Code
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No bone protection medication 1 Calcium and/or vitamin D only 2 Yes - Bisphosphonates, denosumab, romosozumab, teriparatide, raloxifene or HRT (with or without calcium and/or vitamin D) 9 Not known
DD Comments	Calcium or vitamin D includes Calcitriol, calcium and vitamin D or Alpha-calcidol (or one alpha). Bisphosphonates includes: Etidronate, Alendronate, Risedronate, Ibandronate, Zoledronate, Pamidronate. A patient is to be considered as taking osteoporosis specific treatment if: <ul style="list-style-type: none"> • Oral bisphosphonates, prescribed in the last 12 weeks. • Zoledronate, administered in the last 24 months. • Denosumab, administered the last 6 months. • Teriparatide, administered in the last 7 days. • Romosozumab, administered in the last month. These medications may be prescribed with or without calcium and / or vitamin D. For changes to this variable over time, please see current Data Variable Concordance table at https://anzhfr.org/data-access/

Variable Number	7.08
Variable	Re-operation within 120 day follow-up
Variable Name	fop2
Definition	What kind of re-operation has been required (if any) for the patient within 120 days post-surgery?
Justification	To monitor patient outcomes post-surgery
Format	2 digit numeric
Status	Core
Coding Source	Adapted from the UK National Hip Fracture Database
Coding Frame	0 No reoperation 1 Reduction of dislocated prosthesis 2 Washout or debridement 3 Implant removal 4 Revision of internal fixation 5 Conversion to hemiarthroplasty 6 Conversion to total hip replacement 7 Excision arthroplasty 9 Revision arthroplasty 99 Not known
DD Comments	Option 2 washout and debridement includes liner change. Note: record the most significant procedure only.
	The DD comments for this variable have changed over time. Please see current Data Variable Concordance table at https://anzhfr.org/data-access

Variable Number	7.09
Variable	Preliminary date of death
Variable Name	predod
Definition	What was the date of death of the hip fracture patient?
Justification	To monitor patient outcomes and enable reporting of mortality after hip fracture Hip Fracture Clinical Care Standard Indicator 8b.
Format	8 digit, date in DDMMYYYY
Status	Optional, non-core
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 646025). Preliminary Australian date of death obtained from hospital records and/or during 120 day follow-up.
Coding Frame	DD/MM/YYYY
DD Comments	Date of death may be collected either at discharge or during 120-day follow-up. New Zealand date of death may be obtained from the New Zealand Ministry of Health.
	New variable added 1 January 2020. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/

Variable Number	7.10
Variable	.Final date of death
Variable Name	findod
Definition	What was the date of death of the hip fracture patient?
Justification	To monitor patient outcomes and enable reporting of mortality after hip fracture Hip Fracture Clinical Care Standard Indicator 8b.
Format	8 digit, date in DDMMYYYY
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 646025). Final Australian date of death obtained from the National Death Index. New Zealand date of death obtained from the New Zealand Ministry of Health.
Coding Frame	DD/MM/YYYY
DD Comments	Final Australian date of death will be obtained from the National Death Index and final New Zealand date of death will be obtained from the New Zealand Ministry of Health. Date of death will be provided as 30-, 90-, 120- and 365-day mortality flags to facilities/researchers external to Neuroscience Research Australia. New variable added 1 January 2020. Please see current Data Variable Concordance table at https://anzhfr.org/data-access/

Variable Number	7.11
Variable	.Underlying cause of death
Variable Name	undcod
Definition	What was the underlying cause of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian underlying cause of death obtained from the National Death Index. New Zealand underlying cause of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.

Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod1
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
<hr/>	
Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod2
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
<hr/>	
Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod3
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
Variable Number	7.12
Variable	.Other causes of death

Variable Name	Othcod4
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
<hr/>	
Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod5
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
<hr/>	
Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod6
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
Variable Number	7.12
Variable	.Other causes of death
Variable Name	Othcod7
Definition	What were the antecedent causes of death of the hip fracture patient?

Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
Variable Number	7.12
Variable	Other causes of death
Variable Name	Othcod8
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
Variable Number	7.12
Variable	Other causes of death
Variable Name	Othcod9
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.
Variable Number	7.12
Variable	Other causes of death
Variable Name	Othcod10
Definition	What were the antecedent causes of death of the hip fracture patient?
Justification	To enable identification of the underlying cause of death of the hip fracture patient
Format	ANN {.N [N]}

Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian other cause(s) of death obtained from the National Death Index. New Zealand other cause(s) of death obtained from the New Zealand Ministry of Health.
Coding Frame	ICD-10
DD Comments	The disease or injury which initiated the train of morbid events leading directly to a person's death or the circumstances of the incident or violence which produced the fatal injury.

Variable Number	7.13
Variable	.cause of death revision status
Variable Name	cod_revstatus
Definition	What is the revision status of the cause of death?
Justification	All cause of death data from 2006 onwards are subject to a revisions process. Variable included in all data releases
Format	character
Status	Non-core (created centrally)
Coding Source	National Health Data Dictionary, Version 15 (METeOR identifier 307862). Australian underlying cause of death obtained from the National Death Index. New Zealand underlying cause of death obtained from the New Zealand Ministry of Health.
Coding Frame	character
DD Comments	

Variable Number 7.14

Variable **EQ-5D-5L Mobility**

Variable Name Eq5dmob

Definition Please tick the ONE box that best describes your health TODAY

Justification Patient-reported outcome measure

Format 1 digit numeric

Status Core

Coding Source *UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group*

Coding Frame
 1 No problems
 2 Slight problems
 3 Moderate problems
 4 Severe problems
 5 Unable to

DD Comments

Variable Number 7.15

Variable **EQ-5D-5L Self Care**

Variable Name Eq5dcare

Definition Please tick the ONE box that best describes your health TODAY

Justification Patient-reported outcome measure

Format 1 digit numeric

Status Core

Coding Source *UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group*

Coding Frame
 1 No problems
 2 Slight problems
 3 Moderate problems
 4 Severe problems
 5 Unable to

DD Comments

Variable Number 7.16

Variable **EQ-5D-5L Usual Activities**

Variable Name Eq5dact

Definition Please tick the ONE box that best describes your health TODAY

Justification Patient-reported outcome measure

Format 1 digit numeric

Status Core

Coding Source *UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group*

Coding Frame
 1 No problems
 2 Slight problems
 3 Moderate problems
 4 Severe problems
 5 Unable to

DD Comments

Variable Number	7.17
Variable	EQ-5D-5L Pain/Discomfort
Variable Name	Eq5dpain
Definition	Please tick the ONE box that best describes your health TODAY
Justification	Patient-reported outcome measure
Format	1 digit numeric
Status	Core
Coding Source	<i>UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group</i>
Coding Frame	1 = No pain 2 = Slight pain 3 = Moderate pain 4 = Severe pain 5 = Extreme pain
DD Comments	
<hr/>	
Variable Number	7.18
Variable	EQ-5D-5L Anxiety/Depression
Variable Name	Eq5danx
Definition	Please tick the ONE box that best describes your health TODAY
Justification	Patient-reported outcome measure
Format	1 digit numeric
Status	Core
Coding Source	<i>UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group</i>
Coding Frame	1 = Not anxious 2 = Slightly anxious 3 = Moderately anxious 4 = Severely anxious 5 = Extremely anxious
DD Comments	
<hr/>	
Variable Number	7.19
Variable	EQ-5D-5L Health status
Variable Name	Eq5dhealth
Definition	We would like to know how good or bad your health is today. The scale is numbered from 0 to 100. 100 means the best health you can imagine. Zero means the worst health you can imagine.
Justification	Patient-reported outcome measure
Format	2 digit numeric
Status	Core
Coding Source	<i>UK (English) © 2009 EuroQol Group EQ-5D™ is a trade mark of the EuroQol Group</i>
Coding Frame	Likert scale 0-100
DD Comments	

Section 13	52-week follow-up* (50-54 weeks)
-------------------	---

Variable Number	13.01
Variable	Follow-up at 52 weeks
Variable Name	Fup52
Definition	Was the patient followed up at 52 weeks after the index fracture?
Justification	To measure performance against Clinical Standards for Fracture Liaison Services
Format	1 digit numeric
Status	Optional, non-core
Coding Source	Adapted from UK FLS-DB V2.00
Coding Frame	1 Yes 2 No 3 Uncontactable 4 Declined 5 Patient died
DD Comments	This section is only for patients who are recommended bone therapy because of the FLS intervention. Follow up should be at between 48 and 54 weeks after the index fracture (not 52 weeks post assessment). Late follow up - If follow up has been completed, but took place after 54 weeks, please answer 'Yes'. 'No' should only be selected if no follow up is planned.

*52-week follow up is undertaken by the operating hospital

Variable Number	13.02
Variable	.52 Week Follow Up Date
Variable Name	Fup52date
Definition	The date that the "52 week follow up" happened
Justification	To measure performance against Clinical Standards for Fracture Liaison Services
Format	8 digit date, date in DDMMYYYY
Status	Optional, non-core
Coding Source	Adapted from UK FLS-DB V2.00
Coding Frame	dd/mm/yyyy
DD Comments	

Variable Number	13.03
Variable	.52 Week Residence
Variable Name	Fu52residence
Definition	What is the usual place of residence of the patient at the time of the 52 week follow up?
Justification	This enables comparison of the type of accommodation of the person before suffering a fragility fracture with that at follow up assessments. This is an indicator of patient outcome.
Format	1 digit Numeric
Status	Optional, non-core
Coding Source	Adapted from the Australasian Rehabilitation Outcomes Centre Inpatient Dataset, Version 3.0; NSW SNAP Data Collection, Version 4.0
Coding Frame	1 Private residence (including unit in retirement village) 2 Residential aged care facility 3 Other 4 Not done 9 Not known

DD Comments Record the patient's usual accommodation type the time of the 52-week follow up. Residential aged care refers to a supported facility that provides accommodation and care for a person on a long-term basis. This may include multi-purpose services in Australia and private hospitals or rest homes in New Zealand. If the patient lives with a relative or in a community group home or boarding house code 'private residence'. If the patient is in respite care, record their usual place of residence when not in respite care.

Variable Number 13.04
Variable **52 Week Mobility**
Variable Name Walk52fu
Definition The patient's mobility status at the 52-week follow-up
Justification To document the patient's mobility at the time of the 52 week follow up.
Format 1 digit Numeric
Status Optional, non-core
Coding Source Adapted from ANZHFR Data Dictionary V13
Coding Frame 1 Usually walks without walking aids
 2 Usually walks with either a stick or crutch
 3 Usually walks with two aids or frame (with or without assistance of a person)
 4 Usually uses a wheelchair / bed bound
 5 Not done
 9 Not known
DD Comments If a person has different levels of mobility on different surfaces, then record the level of most assistance. For example, inside their residence a person usually walks without a walking aid but when outside the residence the person usually walks with a frame, then the level of mobility recorded is option 3.

Variable Number 13.05
Variable **52 Week Medication**
Variable Name Med52
Definition Did the patient confirm adherence to osteoporosis specific treatment
Justification To document whether the patient was still taking osteoporosis specific treatment
Format 1 digit Numeric
Status Optional, non-core
Coding Source Adapted from UK FLS-DB V2.00
Coding Frame 1 No longer taking osteoporosis specific treatment
 2 Alendronate
 3 Risedronate
 4 Zoledronate
 5 Denosumab
 6 Teriparatide
 7 Testosterone
 8 Systemic Oestrogens
 9 Systemic Oestrogen & Progesterone
 10 Romosozumab
 11 Raloxifene
DD Comments A patient is to be considered as 'on/taking bone protection medication' if:

- For oral-osteoporosis agents patient prescribed in the last 4 weeks.
- For Zoledronate, prescribed in the last 24 months
- For Denosumab, prescribed the last 6 months.
- For Teriparatide, prescribed in the last 7 days.
- For Romosozumab, prescribed in the last month.

Online review of prescriptions may indicate that the patient is taking osteoporosis medication regularly – this is satisfactory. If there is no evidence of this online – patient and / or GP interview will be required

Variable Number	13.06
Variable	Reason for No Medication at 52 Weeks
Variable Name	NoMed52
Definition	What was the reason of the patient not continuing bone protection medication at 52 week follow up?
Justification	To document the reason the patient was no longer taking bone protection medication
Format	1 digit Numeric
Status	Optional, non-core
Coding Source	Adapted from UK FLS-DB V2.00
Coding Frame	1 No longer appropriate (clinician) 2 Informed decline (patient) 3 Side effects 4 Cost to patient 5 Nil obvious 6 Other 7 Not asked 9 Not known
DD Comments	If the patient’s GP or other healthcare professional stops the specific osteoporosis medication for whatever reason (including side effects), please select ‘No longer appropriate (clinician). If the patient stops the medication by the time of the follow up, please select ‘Informed decline (patient)’.

Variable Number	13.07
Variable	Further Falls
Variable Name	furtherfall
Definition	The number of further falls the patient has suffered since the index fracture
Justification	To document the number of further falls since the index fragility fracture suffered by the patient as a measure of patient outcome.
Format	1 digit Numeric
Status	Optional, non-core
Coding Source	
Coding Frame	1 None 2 One 3 Two 4 Three or more 5 Not asked 9 Not known
DD Comments	This is a measure of patient outcome. This is the answer to the question “since the index fracture, have you had any further falls in the last 12 months” or similar.

Variable Number	13.08
Variable	Strength and Balance
Variable Name	SBpartic
Definition	Is the patient still participating in a strength and balance programme?
Justification	To document whether the patient is still participating in strength and balance training.
Format	1 digit Numeric
Status	Optional, non-core
Coding Source	
Coding Frame	1 Yes 2 No 3 Not asked 9 Not known
DD Comments	In the context of this question, a strength and balance programme means that the patient is still carrying out some form of regular activity that aims to improve / maintain their strength and balance. This could be the continuation of an in-home programme that has previously been set or regular attendance at an appropriate community programme. A self-directed programme of regular exercise is also satisfactory.

Variable Number	13.09
Variable	Further Fractures
Variable Name	furtherfract
Definition	Has the patient had a further fragility fracture since the index fracture 52 weeks ago?
Justification	To document whether the patient has had a further fragility fracture since the index fracture 52 weeks ago.
Format	1 digit Numeric
Status	Optional, non-core
Coding Source	
Coding Frame	1 Yes 2 No 3 Not asked 9 Not known